

Introduction to G_2 geometry

Spiro Karigiannis

Department of Pure Mathematics, University of Waterloo

karigiannis@uwaterloo.ca

September 20, 2019

Abstract

These notes give an informal and leisurely introduction to G_2 geometry for beginners. A special emphasis is placed on understanding the special linear algebraic structure in 7 dimensions that is the pointwise model for G_2 geometry, using the octonions. The basics of G_2 -structures are introduced, from a Riemannian geometric point of view, including a discussion of the torsion and its relation to curvature for a general G_2 -structure, as well as the connection to Riemannian holonomy. The history and properties of torsion-free G_2 manifolds are considered, and we stress the similarities and differences with Kähler and Calabi-Yau manifolds. The notes end with a brief survey of three important theorems about compact torsion-free G_2 manifolds.

Contents

1	Aim and scope	2
1.1	History of these notes	2
1.2	Notation	3
2	Motivation	3
3	Algebraic structures from the octonions	5
3.1	Normed division algebras	5
3.2	Induced algebraic structure on $\text{Im } \mathbb{A}$	9
3.3	One-to-one correspondence and classification	12
3.4	Further properties of the cross product in \mathbb{R}^3 and \mathbb{R}^7	13
4	The geometry of G_2-structures	14
4.1	The canonical G_2 -structure on \mathbb{R}^7	14
4.2	G_2 -structures on smooth 7-manifolds	17
4.3	Decomposition of Ω^\bullet into irreducible G_2 representations	19
4.4	The torsion of a G_2 -structure	22
4.5	Relation between curvature and torsion for a G_2 -structure	23
5	Exceptional Riemannian holonomy	24
5.1	Parallel transport and Riemannian holonomy	24
5.2	The Berger classification of Riemannian holonomy	26
6	Torsion-free G_2 manifolds	27
6.1	Irreducible and reducible torsion-free G_2 manifolds	27
6.2	A brief history of irreducible torsion-free G_2 manifolds	28
6.3	Cohomological obstructions to existence in the compact case	29

6.4	Comparison with Kähler and Calabi-Yau manifolds	31
7	Three theorems about compact torsion-free G_2-manifolds	32
7.1	An existence theorem for compact torsion-free G_2 manifolds	32
7.2	The moduli space of compact torsion-free G_2 -structures	33
7.3	A variational characterization of torsion-free G_2 -structures	35
	References	35

1 Aim and scope

The purpose of these lecture notes is to give the reader a gentle introduction to the basic concepts of G_2 geometry, including a brief history of the important early developments of the subject.

At present, there is no “textbook” on G_2 geometry. (This is on the author’s to-do list for the future.) The only references are the classic monograph by Salamon [39] which emphasizes the representation theoretic aspects of Riemannian holonomy, and the book by Joyce [23] which serves as both a text on Kähler and Calabi-Yau geometry as well as a monograph detailing Joyce’s original construction [22] of compact manifolds with G_2 Riemannian holonomy. Both books are excellent resources, but are not easily accessible to beginners. The book by Harvey [17] is at a more appropriate level for new initiates, but is much broader in scope, so it is less focused on G_2 geometry. Moreover, both [39] and [17] predate the important analytic developments that started with Joyce’s seminal contributions.

The aim of the present notes is to attempt to at least psychologically prepare the reader to access the recent literature in the field, which has undergone a veritable explosion in the past few years. The proofs of most of the deeper results are only sketched, with references given to where the reader can find the details, whereas most of the simple algebraic results are proved in detail. Some important aspects of G_2 geometry are unfortunately only briefly mentioned in passing, including the relations to Spin(7)-structures and the intimate connection with spinors and Clifford algebras. Good references for the connection with spinors are Harvey [17], Lawson–Michelsohn [31, Chapter IV. 10], and the more recent paper by Agricola–Chiossi–Friedrich–Höll [2].

These notes are written in a somewhat informal style. In particular, they are meant to be read leisurely. The punchline is sometime spoiled for the benefit of motivation. In addition, results are sometimes explained in more than one way for clarity, and results are not always stated in the correct logical order but rather in an order that (in the humble opinion of the author) is more effective pedagogically. Finally, there is certainly a definite bias towards the personal viewpoint of the author on the subject. In fact, a distinct emphasis is placed on the explicit details of the linear algebraic aspects of G_2 geometry that are consequences of the *nonassociativity of the octonions* \mathbb{O} , as the author believes that this gives good intuition for the striking differences between G_2 -structures and $U(m)$ -structures in general and $SU(m)$ -structures in particular.

The reader is expected to be familiar with graduate level smooth manifold theory and basic Riemannian geometry. Some background in complex and Kähler geometry is helpful, especially to fully appreciate the distinction with G_2 geometry, but is not absolutely essential.

1.1 History of these notes

These lecture notes have been gestating for many years. In their current form, they are a synthesis of lecture notes for several different introductions to G_2 and Spin(7) geometry that have been given by the author at various institutions or workshops over the past decade. Specifically, these are the following, in chronological order:

- October 2006: Seminar; Mathematical Sciences Research Institute; Berkeley.
- November 2008: Seminar series; University of Oxford.
- January/February 2009: Seminar series; University of Waterloo.
- August 2014: ‘ G_2 -manifolds’; Simons Center for Geometry and Physics; Stony Brook.
- September 2014: ‘*Special Geometric Structures in Mathematics and Physics*’; Universität Hamburg.
- August 2017: Minischool on ‘ G_2 -manifolds and related topics’; Fields Institute; Toronto.

The current version of these notes is the first part of the “minischool lectures” on G_2 -geometry collected in the book *Lectures and Surveys on G_2 -geometry and related topics*, published in the *Fields Institute Communications* series by Springer. The other parts of the minischool lectures are

- “Constructions of compact G_2 -holonomy manifolds” by Alexei Kovalev [30]
- “Introduction to G_2 gauge theory” by Thomas Walpuski [41]
- “Geometric flows of G_2 structures” by Jason Lotay [36]
- “Calibrated Submanifolds in G_2 geometry” by Ki Fung Chan and Naichung Conan Leung [34]
- “Calibrated Submanifolds” by Jason Lotay [35]

1.2 Notation

Let (M, g) be a smooth oriented Riemannian n -manifold. We use both vol and μ to denote the Riemannian volume form induced by g and the given orientation. We use the Einstein summation convention throughout. We use $S^2(T^*M)$ to denote the second symmetric power of T^*M .

Given a vector bundle E over M , we use $\Gamma(E)$ to denote the space of smooth sections of E . These spaces are denoted in other ways in the following cases:

- $\Omega^k = \Gamma(\Lambda^k(T^*M))$ is the space of smooth k -forms on M ;
- $\mathcal{S} = \Gamma(S^2(T^*M))$ is the space of smooth symmetric 2-tensors on M .

With respect to the metric g on M , we use \mathcal{S}_0 to denote those sections h of \mathcal{S} that are traceless. That is, \mathcal{S}_0 consists of those sections of \mathcal{S} such that $\text{Tr } h = g^{ij} h_{ij} = 0$ in local coordinates. Then $\mathcal{S} \cong \Omega^0 \oplus \mathcal{S}_0$, where $h \in \mathcal{S}$ is decomposed as $h = \frac{1}{n}(\text{Tr } h)g + h_0$. Then we have $\Gamma(T^*M \otimes T^*M) = \Omega^0 \oplus \mathcal{S}_0 \oplus \Omega^2$, where the splitting is pointwise orthogonal with respect to the metric on $T^*M \otimes T^*M$ induced by g .

Acknowledgements. The author would like to acknowledge Jason Lotay and Naichung Conan Leung for useful discussions on the structuring of these lecture notes. The initial preparation of these notes was done while the author held a Fields Research Fellowship at the Fields Institute. The final preparation of these notes was done while the author was a visiting scholar at the Center of Mathematical Sciences and Applications at Harvard University. The author thanks both the Fields Institute and the CMSA for their hospitality.

2 Motivation

Let (M^n, g) be an n -dimensional smooth Riemannian manifold. For all $p \in M$, we have an n -dimensional real vector space T_pM equipped with a positive-definite inner product g_p , and these “vary smoothly” with $p \in M$. A natural question is the following:

What other “natural structures” can we put on Riemannian manifolds?

What we would like to do is to attach such a “natural structure” to each tangent space T_pM , for all $p \in M$, in a “smoothly varying” way. That is, such a structure corresponds to a smooth section of some

tensor bundle of M , satisfying some natural algebraic condition at each $p \in M$. Let us consider two examples. Let V be an n -dimensional real vector space equipped with a positive-definite inner product g . Note that if we fix an isomorphism $(V, g) \cong (\mathbb{R}^n, \bar{g})$, where \bar{g} is the standard Euclidean inner product on \mathbb{R}^n , then the subgroup of $\text{GL}(n, \mathbb{R})$ preserving this structure is $\text{O}(n)$.

Example 2.1. An *orientation* on V is a nonzero element μ of $\Lambda^n V^*$. Let $\beta = \{e_1, \dots, e_n\}$ be an ordered basis of V . Then $e^1 \wedge \dots \wedge e^n = \lambda \mu$ for some $\lambda \neq 0$. We say that β is positively (respectively, negatively) oriented with respect to μ if $\lambda > 0$ (respectively, $\lambda < 0$). To demand some kind of compatibility with g , we can rescale μ so that $g(\mu, \mu) = 1$. Thus (V, g) admits precisely two orientations. Note that if we fix an isomorphism $(V, g) \cong (\mathbb{R}^n, \bar{g})$, then the subgroup of $\text{O}(n)$ preserving this structures is $\text{SO}(n)$.

On a smooth manifold, an orientation is thus a nowhere-vanishing smooth section μ of $\Lambda^n(T^*M)$. That is, it is a nowhere-vanishing top form. Such a structure *does not always exist*. Specifically, it exists if and only if the real line bundle $\Lambda^n(T^*M)$ is smoothly trivial. In terms of characteristic classes, this condition is equivalent to the vanishing of the first Stiefel-Whitney class $w_1(TM)$ of the tangent bundle. (See [38], for example.) To demand compatibility with g , we can rescale μ by a positive function so that $g_p(\mu_p, \mu_p) = 1$ for all $p \in M$. This normalized μ is the *Riemannian volume form* associated to the metric g and the chosen orientation on M . It is given locally in terms of a positively oriented orthonormal frame $\{e_1, \dots, e_n\}$ of TM by $\mu = e^1 \wedge \dots \wedge e^n$.

An orientation compatible with the metric is called a $\text{SO}(n)$ -structure on M . It is equivalent to a reduction of the structure group of the frame bundle of TM from $\text{GL}(n, \mathbb{R})$ to $\text{SO}(n)$. \blacktriangle

Example 2.2. A *Hermitian structure* on (V, g) is an orthogonal complex structure J . That is, J is a linear endomorphism of V such that $J^2 = -I$ and $g(Jv, Jw) = g(v, w)$ for all $v, w \in V$. It is well-known and easy to check that such a structure exists on V if and only if $n = 2m$ is even. Such a structure allows us to identify the $2m$ -dimensional real vector space V with a m -dimensional complex vector space, where the linear endomorphism J corresponds to multiplication by $\sqrt{-1}$. Note that if we fix an isomorphism $(V, g) \cong (\mathbb{R}^{2m}, \bar{g})$, then the subgroup of $\text{O}(n)$ preserving this structures is $\text{U}(m) = \text{SO}(2m) \cap \text{GL}(m, \mathbb{C})$.

On a Riemannian manifold (M, g) , a Hermitian structure is a smooth section J of the tensor bundle $TM \otimes T^*M = \text{End}(TM)$ such that $J^2 = -I$ (which is called an *almost complex structure*) and such that $g_p(J_p X_p, J_p Y_p) = g_p(X_p, Y_p)$ for all $X_p, Y_p \in T_p M$ (which makes it *orthogonal*). As in Example 2.1, such a structure does not always exist, even if $n = \dim M = 2m$ is even. There are *topological obstructions* to the existence of an almost complex structure, which is equivalent to a reduction of the structure group of the frame bundle of TM from $\text{GL}(2m, \mathbb{R})$ to $\text{GL}(m, \mathbb{C})$. See Massey [37] for discussion on this question.

Further demanding that J be compatible with the metric g (that is, orthogonal) is a reduction of the structure group of the frame bundle of TM from $\text{GL}(2m, \mathbb{R})$ to $\text{U}(m)$. For this reason a Hermitian structure on M^{2m} is sometimes also called a $\text{U}(m)$ -structure. Readers can consult [12] for a comprehensive treatment of the geometry of general $\text{U}(m)$ -structures. \blacktriangle

Again, let V be an n -dimensional real vector space equipped with a positive-definite inner product g . A G_2 -structure is a special algebraic structure we can consider on (V, g) only when $n = 7$. In this case, if we fix an isomorphism $(V, g) \cong (\mathbb{R}^7, \bar{g})$, then G_2 is the subgroup of $\text{SO}(7)$ preserving this special algebraic structure. In order to describe this structure at the level of linear algebra, we first need to discuss the *octonions*, which we do in Section 3. Then G_2 -structures are defined and studied in Section 4. For the purposes of this motivational section, all the reader needs to know is that a “ G_2 -structure” corresponds to a special kind of 3-form φ on M^7 .

Suppose we have a “natural structure” on a Riemannian manifold (M^n, g) , such as that of Examples 2.1 or 2.2 or the mysterious G_2 -structure that is the subject of the present notes. Since we have a Riemannian metric g , we have a Levi-Civita connection ∇ and we can further ask for the “natural structure” to be *parallel* or *covariantly constant* with respect to ∇ . For example:

- If μ is an orientation (Riemannian volume form) on (M^n, g) , then it is *always* parallel.

- If J is an orthogonal almost complex structure on (M^{2m}, g) , then if we have $\nabla J = 0$, we say that (M, g, J) is a *Kähler* manifold. Such manifolds have been classically well-studied.
- If φ is a G_2 -structure on (M^7, g) , then if we have $\nabla\varphi = 0$, we say that (M, g, φ) is a *torsion-free G_2 manifold*. Such manifolds are discussed in Section 6 of the present notes.

3 Algebraic structures from the octonions

In this section we give an introduction to the algebra of the *octonions* \mathbb{O} , an 8-dimensional real normed division algebra, and to the induced algebraic structure on $\text{Im } \mathbb{O}$, the 7-dimensional space of *imaginary octonions*. We do this by discussing both normed division algebras and spaces equipped with a cross product, and then relating the two concepts. This is not strictly necessary if the intent is to simply consider G_2 -structures, but it has the pedagogical benefit of putting both G_2 and $\text{Spin}(7)$ geometry into the proper wider context of geometries associated to real normed division algebras. (See Leung [32] for more on this perspective.)

We do not discuss all of the details here, but we do prove many of the important simple results. More details on the algebraic structure of the octonions can be found in Harvey [17], Harvey–Lawson [18], and Salamon–Walpuski [40], for example.

3.1 Normed division algebras

Let $\mathbb{A} = \mathbb{R}^n$ be equipped with the standard Euclidean inner product $\langle \cdot, \cdot \rangle$.

Definition 3.1. We say that \mathbb{A} is a *normed division algebra* if \mathbb{A} has the structure of a (*not necessarily associative!*) algebra over \mathbb{R} with multiplicative identity $1 \neq 0$ such that

$$\|ab\| = \|a\| \|b\| \quad \text{for all } a, b \in \mathbb{A}, \quad (3.2)$$

where $\|a\|^2 = \langle a, a \rangle$ is the usual Euclidean norm on \mathbb{R}^n induced from $\langle \cdot, \cdot \rangle$. Equation (3.2) imposes a *compatibility condition* between the inner product and the algebra structure on \mathbb{A} . ▲

Remark 3.3. This is not the most general definition possible, but it suffices for our purposes. See [18, Appendices IV.A and IV.B] for more details. ▲

We discuss examples of normed division algebras later in this section, although it is clear that the standard algebraic structures on \mathbb{R} and $\mathbb{C} \cong \mathbb{R}^2$ give examples. We now define some additional structures and investigate some properties of normed division algebras. It is truly remarkable how many far reaching consequences arise solely from the fundamental identity (3.2).

Definition 3.4. Let \mathbb{A} be a normed division algebra. Define the *real part* of \mathbb{A} , denoted $\text{Re } \mathbb{A}$, to be the span over \mathbb{R} of the multiplicative identity $1 \in \mathbb{A}$. That is, $\text{Re } \mathbb{A} = \{t1 : t \in \mathbb{R}\}$. Define the *imaginary part* of \mathbb{A} , denoted $\text{Im } \mathbb{A}$, to be the orthogonal complement of $\text{Re } \mathbb{A}$ with respect to the Euclidean inner product on $\mathbb{A} = \mathbb{R}^n$. That is, $\text{Im } \mathbb{A} = (\text{Re } \mathbb{A})^\perp \cong \mathbb{R}^{n-1}$. Given $a \in \mathbb{A}$, there exists a unique decomposition

$$a = \text{Re } a + \text{Im } a, \quad \text{where } \text{Re } a \in \text{Re } \mathbb{A} \text{ and } \text{Im } a \in \text{Im } \mathbb{A}.$$

We define the *conjugate* \bar{a} of a to be the element

$$\bar{a} = \text{Re } a - \text{Im } a. \quad \text{▲}$$

Note that the map $a \mapsto \bar{a}$ is a linear involution of \mathbb{A} , and is precisely the isometry that is minus the reflection across the hyperplane $\text{Im } \mathbb{A}$ of \mathbb{A} . It is clear that

$$\text{Re } a = \frac{1}{2}(a + \bar{a}) \quad \text{and} \quad \text{Im } a = \frac{1}{2}(a - \bar{a}). \quad (3.5)$$

As a result, we deduce that

$$\bar{a} = -a \quad \text{if and only if} \quad a \in \text{Im } \mathbb{A}. \quad (3.6)$$

We now derive a slew of important identities that are all consequences of the defining property (3.2).

Lemma 3.7. *Let $a, b, c \in \mathbb{A}$. Then we have*

$$\langle ac, bc \rangle = \langle ca, cb \rangle = \langle a, b \rangle \|c\|^2, \quad (3.8)$$

and

$$\langle a, bc \rangle = \langle a\bar{c}, b \rangle, \quad \langle a, cb \rangle = \langle \bar{c}a, b \rangle. \quad (3.9)$$

Moreover, we also have

$$\overline{ab} = \bar{b}\bar{a}. \quad (3.10)$$

Proof. First observe that

$$\begin{aligned} \|(a+b)c\|^2 &= \|ac+bc\|^2 = \|ac\|^2 + 2\langle ac, bc \rangle + \|bc\|^2, \\ \|a+b\|^2 \|c\|^2 &= (\|a\|^2 + 2\langle a, b \rangle + \|b\|^2) \|c\|^2. \end{aligned}$$

Equating the left hand sides above using the fundamental identity (3.2), and again using (3.2) to cancel the corresponding first and third terms on the right hand sides, we obtain

$$\langle ac, bc \rangle = \langle a, b \rangle \|c\|^2. \quad (3.11)$$

Similarly we can show

$$\langle ca, cb \rangle = \langle a, b \rangle \|c\|^2. \quad (3.12)$$

Thus we have established (3.8). Consider the first equation in (3.9). It is clearly satisfied when c is real, since then $\bar{c} = c$ and the inner product $\langle \cdot, \cdot \rangle$ is bilinear over \mathbb{R} . Because both sides of the equation are linear in c , it is enough to consider the situation when c is purely imaginary, in which case $\bar{c} = -c$. Then c is orthogonal to 1, so $\|1+c\|^2 = 1 + \|c\|^2$. Applying (3.2) and (3.8), we find

$$\begin{aligned} \langle a, b \rangle (1 + \|c\|^2) &= \langle a, b \rangle \|1+c\|^2 = \langle a(1+c), b(1+c) \rangle \\ &= \langle a+ac, b+bc \rangle = \langle a, b \rangle + \langle ac, bc \rangle + \langle a, bc \rangle + \langle ac, b \rangle \\ &= \langle a, b \rangle + \langle a, b \rangle \|c\|^2 + \langle a, bc \rangle + \langle ac, b \rangle. \end{aligned}$$

Thus we have $\langle a, bc \rangle = -\langle ac, b \rangle = \langle a\bar{c}, b \rangle$. This establishes the first equation in (3.9). The other is proved similarly. Using (3.9) and the fact that conjugation is an isometry, we have

$$\langle \bar{ab}, c \rangle = \langle ab, \bar{c} \rangle = \langle b, \bar{a}\bar{c} \rangle = \langle bc, \bar{a} \rangle = \langle c, \bar{b}\bar{a} \rangle.$$

Since this holds for all $c \in \mathbb{A}$, we deduce that $\overline{ab} = \bar{b}\bar{a}$. □

Lemma 3.7 has several important corollaries.

Corollary 3.13. *Let $a, b, c \in \mathbb{A}$. Then we have*

$$a(\bar{b}c) + b(\bar{a}c) = 2\langle a, b \rangle c, \quad (3.14)$$

$$(\bar{a}b)c + (\bar{a}\bar{c})b = 2\langle b, c \rangle a, \quad (3.15)$$

$$\bar{a}b + b\bar{a} = 2\langle a, b \rangle 1. \quad (3.16)$$

Proof. Polarizing (3.8), we have

$$\begin{aligned} \langle a, b \rangle \|c + d\|^2 &= \langle a(c + d), b(c + d) \rangle \\ \langle a, b \rangle (\|c\|^2 + 2\langle c, d \rangle + \|d\|^2) &= \langle ac, bc \rangle + \langle ad, bc \rangle + \langle ac, bd \rangle + \langle ad, bd \rangle, \end{aligned}$$

and hence upon using (3.8) to cancel the corresponding first and last terms on each side, we get

$$\langle ad, bc \rangle + \langle ac, bd \rangle = 2\langle a, b \rangle \langle c, d \rangle. \quad (3.17)$$

Using (3.9), we can write the above as

$$\langle d, \bar{a}(bc) \rangle + \langle \bar{b}(ac), d \rangle = 2\langle a, b \rangle \langle c, d \rangle.$$

Since the above holds for any $d \in \mathbb{A}$, we deduce that

$$\bar{a}(bc) + \bar{b}(ac) = 2\langle a, b \rangle c.$$

Replacing $a \mapsto \bar{a}$ and $b \mapsto \bar{b}$ and using the fact that conjugation is an isometry, we obtain (3.14). Equation (3.15) is obtained similarly. Alternatively, one can take the conjugate of (3.14) and use the relation (3.10). Finally, (3.16) is the special case of (3.14) when $c = 1$. \square

Corollary 3.18. *Let $a, b, c \in \text{Im } \mathbb{A}$. Then we have*

$$a(bc) + b(ac) = -2\langle a, b \rangle c, \quad (3.19)$$

$$(ab)c + (ac)b = -2\langle b, c \rangle a, \quad (3.20)$$

$$ab + ba = -2\langle a, b \rangle 1. \quad (3.21)$$

Proof. These are immediate from Corollary 3.13 and equation (3.6). \square

Corollary 3.22. *Let $a, b \in \mathbb{A}$. Then we have*

$$\langle a, b \rangle = \text{Re}(a\bar{b}) = \text{Re}(b\bar{a}) = \text{Re}(\bar{b}a) = \text{Re}(\bar{a}b) \quad (3.23)$$

and

$$\|a\|^2 = a\bar{a} = \bar{a}a. \quad (3.24)$$

Proof. Using (3.9), we have $\langle a, b \rangle = \langle a, b1 \rangle = \langle a\bar{b}, 1 \rangle = \text{Re}(a\bar{b})$. The remaining equalities in (3.23) follow from the symmetry of $\langle \cdot, \cdot \rangle$ and the fact that conjugation is an isometry. From (3.10), we find $\bar{\bar{a}a} = \bar{a}\bar{\bar{a}} = \bar{a}a$, so $\bar{a}a$ is real. Equation (3.24) thus follows from (3.23). \square

Corollary 3.25. *Let $a \in \mathbb{A}$. Then $a^2 = aa$ is real if and only if a is either real or imaginary.*

Proof. Write $a = \text{Re } a + \text{Im } a$. Since $\overline{\text{Im } a} = -\text{Im } a$, from (3.24) we have $(\text{Im } a)^2 = -(\text{Im } a)(\overline{\text{Im } a}) = -\|\text{Im } a\|^2$. Thus we have

$$a^2 = (\text{Re } a + \text{Im } a)(\text{Re } a + \text{Im } a) = ((\text{Re } a)^2 - \|\text{Im } a\|^2)1 + 2(\text{Re } a)(\text{Im } a).$$

Since the first term on the right hand side above is always real and the second term is always imaginary, we conclude that a^2 is real if and only if $(\text{Re } a)(\text{Im } a) = 0$, which means that either $\text{Re } a = 0$ or $\text{Im } a = 0$. \square

Corollary 3.26. *Let $a, c \in \mathbb{A}$. Then we have*

$$\begin{aligned} (ac)\bar{c} &= a(c\bar{c}) = \|c\|^2 a = a(\bar{c}c) = (a\bar{c})c, \\ a(\bar{a}c) &= (a\bar{a})c = \|a\|^2 c = (\bar{a}a)c = \bar{a}(ac). \end{aligned} \quad (3.27)$$

Proof. Observe from (3.9) and (3.8) that

$$\langle (ac)\bar{c}, b \rangle = \langle ac, bc \rangle = \langle a, b \rangle \|c\|^2 = \langle a \|c\|^2, b \rangle = \langle a(c\bar{c}), b \rangle.$$

Since this holds for all $b \in \mathbb{A}$, we deduce that

$$(ac)\bar{c} = a(c\bar{c}).$$

The rest of the first identity in (3.27) follows by interchanging c and \bar{c} . The second identity in (3.27) is proved similarly. \square

We now introduce two fundamental \mathbb{A} -valued multilinear maps on \mathbb{A} .

Definition 3.28. Let \mathbb{A} be a normed division algebra. Define a bilinear map $[\cdot, \cdot] : \mathbb{A}^2 \rightarrow \mathbb{A}$ by

$$[a, b] = ab - ba \quad \text{for all } a, b \in \mathbb{A}. \quad (3.29)$$

The map $[\cdot, \cdot]$ is called the *commutator* of \mathbb{A} .

Define a trilinear map $[\cdot, \cdot, \cdot] : \mathbb{A}^3 \rightarrow \mathbb{A}$ by

$$[a, b, c] = (ab)c - a(bc) \quad \text{for all } a, b, c \in \mathbb{A}. \quad (3.30)$$

The map $[\cdot, \cdot, \cdot]$ is called the *associator* of \mathbb{A} . \blacktriangle

It is clear that the commutator vanishes identically on \mathbb{A} if and only if \mathbb{A} is commutative, and similarly the associator vanishes identically on \mathbb{A} if and only if \mathbb{A} is associative.

Proposition 3.31. *The commutator and associator are both alternating. That is, they are totally skew-symmetric in their arguments.*

Proof. The commutator is clearly alternating. Because \mathbb{A} is an algebra over \mathbb{R} , the associator clearly vanishes if any of the arguments are purely real. Thus, because the associator is trilinear it suffices to show that \mathbb{A} is alternating on $(\text{Im } \mathbb{A})^3$. If $a, b \in \text{Im } \mathbb{A}$, then $\bar{a} = -a$ and $\bar{b} = -b$. Thus by (3.27) we find that

$$-[a, a, b] = [a, \bar{a}, b] = (a\bar{a})b - a(\bar{a}b) = 0.$$

Similarly we have $-[a, b, b] = [a, \bar{b}, b] = (a\bar{b})b - a(\bar{b}b) = 0$. Thus $[\cdot, \cdot, \cdot]$ is alternating in its first two arguments and in its last two arguments. Thus $[a, b, a] = -[a, a, b] = 0$ as well. \square

The next result says that both the commutator and the associator restrict to vector-valued alternating multilinear forms on $\text{Im } \mathbb{A}$.

Lemma 3.32. *If $a, b, c \in \text{Im } \mathbb{A}$, then $[a, b] \in \text{Im } \mathbb{A}$ and $[a, b, c] \in \text{Im } \mathbb{A}$.*

Proof. We need to show that $[a, b]$ and $[a, b, c]$ are orthogonal to 1. Using the fact that $\bar{a} = -a$ for any $a \in \text{Im } \mathbb{A}$, and the identities (3.9) and (3.46), we compute

$$\begin{aligned} \langle [a, b], 1 \rangle &= \langle ab - ba, 1 \rangle = \langle b, \bar{a} \rangle - \langle a, \bar{b} \rangle \\ &= -\langle b, a \rangle + \langle a, b \rangle = 0. \end{aligned}$$

Similarly, noting that $\overline{bc} = \bar{c}\bar{b} = (-c)(-b) = cb$, we have

$$\begin{aligned} \langle [a, b, c], 1 \rangle &= \langle (ab)c - a(bc), 1 \rangle = \langle ab, \bar{c} \rangle - \langle bc, \bar{a} \rangle \\ &= -\langle ab, c \rangle + \langle bc, a \rangle = -\langle a, \bar{c}b \rangle + \langle bc, a \rangle \\ &= \langle a, cb + bc \rangle = \langle a, bc + \overline{bc} \rangle = 2\langle a, \text{Re}(bc) \rangle = 0, \end{aligned}$$

as claimed. \square

Proposition 3.33. *Let $a, b, c, d \in \mathbb{A}$. The multilinear expressions $\langle a, [b, c] \rangle$ and $\langle a, [b, c, d] \rangle$ are both totally skew-symmetric in their arguments.*

Proof. The commutator and the associator are alternating by Proposition 3.31. Thus we need only show that $\langle a, [a, b] \rangle = 0$ and $\langle a, [a, b, c] \rangle = 0$. Using the identity (3.8) we compute

$$\langle a, [a, b] \rangle = \langle a, ab - ba \rangle = \|a\|^2 \langle 1, b \rangle - \|a\|^2 \langle 1, b \rangle = 0,$$

and similarly using (3.8) and (3.9) we have

$$\begin{aligned} \langle a, [a, b, c] \rangle &= \langle a, (ab)c - a(bc) \rangle = \langle a\bar{c}, ab \rangle - \|a\|^2 \langle 1, bc \rangle \\ &= \|a\|^2 \langle \bar{c}, b \rangle - \|a\|^2 \langle \bar{c}, b \rangle = 0 \end{aligned}$$

as claimed. \square

3.2 Induced algebraic structure on $\text{Im } \mathbb{A}$

Let $\mathbb{A} \cong \mathbb{R}^n$ be a normed division algebra with imaginary part $\text{Im } \mathbb{A} \cong \mathbb{R}^{n-1}$. We define several objects on $\text{Im } \mathbb{A}$ induced from the algebra structure on \mathbb{A} . Motivated by Lemma 3.32 and Proposition 3.33 the following definition is natural. The factor of $\frac{1}{2}$ is for convenience, to avoid factors of 2 in equations (3.45) and (3.63).

Definition 3.34. Recall the statement of Proposition 3.33. Define a 3-form φ and a 4-form ψ on $\text{Im } \mathbb{A}$ as follows:

$$\varphi(a, b, c) = \frac{1}{2} \langle a, [b, c] \rangle = \frac{1}{2} \langle [a, b], c \rangle \quad \text{for } a, b, c \in \text{Im } \mathbb{A}, \quad (3.35)$$

$$\psi(a, b, c, d) = \frac{1}{2} \langle a, [b, c, d] \rangle = -\frac{1}{2} \langle [a, b, c], d \rangle \quad \text{for } a, b, c, d \in \text{Im } \mathbb{A}. \quad (3.36)$$

The form $\varphi \in \Lambda^3(\text{Im } \mathbb{A})^*$ is called the *associative* 3-form, and the form $\psi \in \Lambda^4(\text{Im } \mathbb{A})^*$ is called the *coassociative* 4-form for reasons that become clear in the context of calibrated geometry [35, 34]. \blacktriangle

The 3-form φ is intimately related to another algebraic operation on $\text{Im } \mathbb{A}$ that is fundamental in G_2 -geometry, given by the following definition.

Definition 3.37. Define a bilinear map $\times : \mathbb{A}^2 \rightarrow \mathbb{A}$ by

$$a \times b = \text{Im}(ab) \quad \text{for all } a, b \in \text{Im } \mathbb{A}. \quad (3.38)$$

Essentially, since the product in \mathbb{A} of two imaginary elements need not be imaginary, we project to the imaginary part to define \times . The bilinear map \times is called the *vector cross product* on $\text{Im } \mathbb{A}$ induced by the algebraic structure on \mathbb{A} . \blacktriangle

The vector cross product \times has several interesting properties.

Lemma 3.39. *Let $a, b \in \text{Im } \mathbb{A}$. Then we have*

$$a \times b = -b \times a, \quad (3.40)$$

$$\langle a \times b, a \rangle = 0, \quad \text{so } (a \times b) \perp a \text{ and } (a \times b) \perp b, \quad (3.41)$$

$$\text{Re}(ab) = -\langle a, b \rangle 1. \quad (3.42)$$

Proof. Recall that $\bar{a} = -a$ and $\bar{b} = -b$. Thus from (3.5) and (3.10), we have

$$2a \times b = ab - \overline{ab} = ab - ba = [a, b]. \quad (3.43)$$

Thus $a \times b = -b \times a$.

Since $a \in \text{Im } \mathbb{A}$, equation (3.38) show that $\langle a \times b, a \rangle = \langle \text{Im}(ab), a \rangle = \langle ab, a \rangle$. Thus, using (3.8) we get $\langle a \times b, a \rangle = \langle ab, a \rangle = \langle ab, a1 \rangle = \|a\|^2 \langle b, 1 \rangle = 0$ because $b \in \text{Im } \mathbb{A}$ is orthogonal to $1 \in \text{Re } \mathbb{A}$.

Since $\bar{b} = -b$, equation (3.23) gives $\langle a, b \rangle = \text{Re}(a\bar{b}) = -\text{Re}(ab)$, which is (3.42). \square

Combining equations (3.42) and (3.38) gives us that the decomposition of $ab \in \mathbb{A}$ into real and imaginary parts is given by

$$ab = -\langle a, b \rangle 1 + a \times b. \quad (3.44)$$

It then follows from (3.43) and (3.35) that

$$\varphi(a, b, c) = \langle a \times b, c \rangle \quad \text{for } a, b, c \in \text{Im } \mathbb{A}. \quad (3.45)$$

Note that since $a \times b - ab$ is real by (3.44), we can equivalently write (3.45) as

$$\varphi(a, b, c) = \langle ab, c \rangle \quad \text{for } a, b, c \in \text{Im } \mathbb{A}. \quad (3.46)$$

Lemma 3.47. *Let $a, b, c \in \text{Im } \mathbb{A}$. Then we have*

$$a(bc) = -\frac{1}{2}[a, b, c] - \varphi(a, b, c)1 - \langle b, c \rangle a + \langle a, c \rangle b - \langle a, b \rangle c. \quad (3.48)$$

Proof. Applying the three identities in Corollary 3.18 repeatedly, we compute

$$\begin{aligned} a(bc) &= -b(ac) - 2\langle a, b \rangle c \\ &= -b(-ca - 2\langle a, c \rangle 1) - 2\langle a, b \rangle c \\ &= b(ca) + 2\langle a, c \rangle b - 2\langle a, b \rangle c \\ &= -c(ba) - 2\langle b, c \rangle a + 2\langle a, c \rangle b - 2\langle a, b \rangle c. \end{aligned}$$

Also, putting $c \mapsto c$ and $b \mapsto ab$ in (3.16) and using (3.46) gives

$$c(ba) - (ab)c = c(\overline{ab}) + (ab)\bar{c} = 2\langle ab, c \rangle 1 = 2\varphi(a, b, c)1.$$

Combining the above two expressions gives

$$a(bc) = -(ab)c - 2\varphi(a, b, c)1 - 2\langle b, c \rangle a + 2\langle a, c \rangle b - 2\langle a, b \rangle c.$$

Using $[a, b, c] = (ab)c - a(bc)$ to eliminate $(ab)c$ above and rearranging gives (3.48). \square

Equation (3.48) is used to establish the following two corollaries.

Corollary 3.49. *Let $a, b, c \in \text{Im } \mathbb{A}$. The vector cross product \times on $\text{Im } \mathbb{A}$ satisfies*

$$\|a \times b\|^2 = \|a\|^2 \|b\|^2 - \langle a, b \rangle^2 = \|a \wedge b\|^2, \quad (3.50)$$

and

$$a \times (b \times c) = -\langle a, b \rangle c + \langle a, c \rangle b - \frac{1}{2}[a, b, c]. \quad (3.51)$$

Proof. Let $a, b \in \text{Im } \mathbb{A}$. Using (3.44) we have $ab = -\langle a, b \rangle 1 + a \times b$ and $ba = -\langle b, a \rangle 1 + b \times a = -\langle a, b \rangle - a \times b$. Thus we have

$$\langle ab, ba \rangle = \langle -\langle a, b \rangle 1 + a \times b, -\langle a, b \rangle 1 - a \times b \rangle = \langle a, b \rangle^2 - \|a \times b\|^2.$$

Using the above expression and equations (3.43) and (3.2), we compute

$$\begin{aligned} 4\|a \times b\|^2 &= \langle ab - ba, ab - ba \rangle = \|ab\|^2 + \|ba\|^2 - 2\langle ab, ba \rangle \\ &= \|a\|^2 \|b\|^2 + \|b\|^2 \|a\|^2 - 2(\langle a, b \rangle^2 - \|a \times b\|^2), \end{aligned}$$

which simplifies to (3.50). Again using (3.44) twice and (3.45) we compute

$$\begin{aligned} a \times (b \times c) &= \langle a, b \times c \rangle 1 + a(b \times c) \\ &= \varphi(a, b, c)1 + a(\langle b, c \rangle + bc) \\ &= a(bc) + \varphi(a, b, c)1 + \langle b, c \rangle a. \end{aligned}$$

Substituting (3.48) for $a(bc)$ above gives (3.51). \square

Corollary 3.52. *Let $a, b, c, d \in \text{Im } \mathbb{A}$. Then the following identity holds:*

$$\|\tfrac{1}{2}[a, b, c]\|^2 + (\varphi(a, b, c))^2 = \|a \wedge b \wedge c\|^2. \quad (3.53)$$

Proof. Recall from Lemma 3.32 and Proposition 3.33 that $[a, b, c]$ is purely imaginary and is orthogonal to a, b, c . Thus, taking the norm squared of (3.48) and using the fundamental relation (3.2), we have

$$\begin{aligned} \|a\|^2 \|b\|^2 \|c\|^2 &= \|a\|^2 \|bc\|^2 = \|a(bc)\|^2 \\ &= \|\tfrac{1}{2}[a, b, c]\|^2 + (\varphi(a, b, c))^2 + \|a\|^2 \langle b, c \rangle^2 + \|b\|^2 \langle a, c \rangle^2 + \|c\|^2 \langle a, b \rangle^2 \\ &\quad - 2\langle b, c \rangle \langle a, c \rangle \langle a, b \rangle + 2\langle b, c \rangle \langle a, b \rangle \langle a, c \rangle - 2\langle a, c \rangle \langle a, b \rangle \langle b, c \rangle. \end{aligned}$$

This can be rearranged to yield

$$\begin{aligned} \|\tfrac{1}{2}[a, b, c]\|^2 + (\varphi(a, b, c))^2 &= \|a\|^2 \|b\|^2 \|c\|^2 - \|a\|^2 \langle b, c \rangle^2 - \|b\|^2 \langle a, c \rangle^2 \\ &\quad - \|c\|^2 \langle a, b \rangle^2 + 2\langle a, b \rangle \langle a, c \rangle \langle b, c \rangle, \end{aligned}$$

which is precisely (3.53). \square

Remark 3.54. Comparing equations (3.35) and (3.45), one is tempted from (3.36) to think of the expression $\tfrac{1}{2}[a, b, c]$ as some kind of 3-fold vector cross product $P(a, b, c)$, as it is a trilinear vector valued alternating form on $\text{Im } \mathbb{A}$. However, equation (3.53) says that $\|a \wedge b \wedge c\|^2 - \|P(a, b, c)\|^2$ is nonzero in general, whereas (3.50) says that $\|a \wedge b\|^2 - \|a \times b\|^2 = 0$ always. There *is* a notion of 3-fold vector cross product (see Remark 3.59 below) but $[\cdot, \cdot, \cdot]$ on $\text{Im } \mathbb{A}$ is not one of them. In fact, equation (3.53) is the *calibration inequality* for φ . It says that $|\varphi(a, b, c)| \leq 1$ whenever a, b, c are orthonormal, with equality if and only if $[a, b, c] = 0$. See [35, 34] in the present volume for more about the aspects of G_2 geometry related to *calibrations*. \blacktriangle

Equations (3.41) and (3.50) for the vector cross product \times induced from the algebraic structure on \mathbb{A} motivate the following general definition.

Definition 3.55. Let $\mathbb{V} = \mathbb{R}^m$, equipped with the usual Euclidean inner product. We say that V has a *vector cross product*, which we usually simply call a *cross product*, if there exists a *skew-symmetric* bilinear map $\times : \mathbb{V}^2 \rightarrow \mathbb{V}$ such that, for all $a, b, c \in \mathbb{V}$, we have

$$\langle a \times b, a \rangle = 0, \quad \text{so } (a \times b) \perp a \text{ and } (a \times b) \perp b, \quad (3.56)$$

$$\|a \times b\|^2 = \|a\|^2 \|b\|^2 - \langle a, b \rangle^2 = \|a \wedge b\|^2. \quad (3.57)$$

Note that (3.56) and (3.57) are precisely (3.41) and (3.50), respectively. \blacktriangle

Remark 3.58. The fact that \times is skew-symmetric and bilinear is equivalent to saying that \times is a *linear map*

$$\times : \Lambda^2(\mathbb{V}) \rightarrow \mathbb{V}.$$

Then (3.57) says that \times is *length preserving on decomposable elements of $\Lambda^2(\mathbb{V})$* . \blacktriangle

Remark 3.59. In Definition 3.55 we have really defined a special class of vector cross product, called a *2-fold vector cross product*. A more general notion of *k-fold vector cross product* [3] exists. When $k = 1$ such a structure is an orthogonal complex structure. When $k = 3$ such a structure is related to Spin(7)-geometry. See also Lee–Leung [33] for more details. Another extensive recent reference for general vector cross products is Cheng–Karigiannis–Madnick [7, Section 2]. \blacktriangle

We have seen that any normed division algebra \mathbb{A} gives a vector cross product on $\mathbb{V} = \text{Im } \mathbb{A}$. In the next section we show that we can also go the other way.

3.3 One-to-one correspondence and classification

We claim that the normed division algebras are in *one-to-one correspondence* with the spaces admitting cross products. The correspondence is seen explicitly as follows. Let \mathbb{A} be a normed division algebra. In Section 3.2 we showed that $\mathbb{V} = \text{Im } \mathbb{A}$ has a cross product. Conversely, suppose $\mathbb{V} = \mathbb{R}^m$ has a cross product \times . Define $\mathbb{A} = \mathbb{R} \oplus \mathbb{V} = \mathbb{R}^{m+1}$, with the Euclidean inner product. That is,

$$\langle (s, v), (t, w) \rangle = st + \langle v, w \rangle \quad \text{for } s, t \in \mathbb{R} \text{ and } v, w \in \mathbb{V}.$$

Define a multiplication on \mathbb{A} by

$$(s, v)(t, w) = (st - \langle v, w \rangle, sw + tv + v \times w). \quad (3.60)$$

The multiplication defined in (3.60) is clearly bilinear over \mathbb{R} , so it gives \mathbb{A} the structure of a (not necessarily associative) algebra over \mathbb{R} . It is also clear from (3.60) that $(1, 0)$ is a multiplicative identity on \mathbb{A} . We need to check that (3.2) is satisfied. We compute:

$$\begin{aligned} \|(s, v)(t, w)\|^2 &= (st - \langle v, w \rangle)^2 + \|sw + tv + v \times w\|^2 \\ &= s^2 t^2 - 2st \langle v, w \rangle + \langle v, w \rangle^2 + s^2 \|w\|^2 + t^2 \|v\|^2 + \|v \times w\|^2 \\ &\quad + 2st \langle v, w \rangle + 2s \langle w, v \times w \rangle + 2t \langle v, v \times w \rangle. \end{aligned}$$

Using (3.51) and (3.56), the above expression simplifies to

$$\begin{aligned} \|(s, v)(t, w)\|^2 &= s^2 t^2 + s^2 \|w\|^2 + t^2 \|v\|^2 + \|v\|^2 \|w\|^2 \\ &= (s^2 + \|v\|^2)(t^2 + \|w\|^2) = \|(s, v)\| \|(t, w)\|, \end{aligned}$$

verifying (3.2).

Normed division algebras were classified by Hurwitz in 1898. A proof using the *Cayley–Dickson doubling construction* can be found in [17, Chapter 6] or [18, Appendix IV.A]. There are exactly four possibilities, up to isomorphism.

The four normed division algebras are given by the following table:

$n = \dim \mathbb{A}$	1	2	4	8
Symbol	\mathbb{R}	$\mathbb{C} \cong \mathbb{R}^2$	$\mathbb{H} \cong \mathbb{R}^4$	$\mathbb{O} \cong \mathbb{R}^8$
Name	Real numbers	Complex numbers	Quaternions or Hamilton numbers	Octonions or Cayley numbers

Each algebra in the above table is a *subalgebra* of the next one. In particular, the multiplicative unit in all cases is the usual multiplicative identity $1 \in \mathbb{R}$. Moreover, as we enlarge the algebras $\mathbb{R} \rightarrow \mathbb{C} \rightarrow \mathbb{H} \rightarrow \mathbb{O}$, we lose some algebraic property at each step. From \mathbb{R} to \mathbb{C} we lose the *natural ordering*. From \mathbb{C} to \mathbb{H} we lose *commutativity*. And from \mathbb{H} to \mathbb{O} we lose *associativity*.

The octonions \mathbb{O} are also called the *exceptional* division algebra and the geometries associated to \mathbb{O} are known as *exceptional geometries*.

By the one-to-one correspondence between normed division algebras and spaces admitting cross products, we deduce that there exist precisely four spaces with cross product, given by the following table:

$m = \dim \mathbb{V}$	0	1	3	7
Symbol	$\{0\} \cong \text{Im } \mathbb{R}$	$\mathbb{R} \cong \text{Im } \mathbb{C}$	$\mathbb{R}^3 \cong \text{Im } \mathbb{H}$	$\mathbb{R}^7 \cong \text{Im } \mathbb{O}$
Cross product \times	trivial	trivial	standard (Hodge star)	exceptional
the 3-form $\varphi \in \Lambda^3(\mathbb{V}^*)$	0	0	$\varphi = \mu$ is the standard volume form	φ is the associative 3-form
the $(m-3)$ -form $\star\varphi \in \Lambda^{m-3}(\mathbb{V}^*)$	0	0	$\star\varphi = 1 \in \Lambda^0(\mathbb{V}^*) \cong \mathbb{R}$ is the multiplicative unit	$\star\varphi = \psi \in \Lambda^4(\mathbb{V}^*)$ is the coassociative 4-form

Remark 3.61. Here are some remarks concerning the above table:

- (i) When $m = 0, 1$, the cross product \times is trivial because $\Lambda^2(\mathbb{V}) = \{0\}$ in these cases.
- (ii) When $m = 3$ we recover the standard cross product on \mathbb{R}^3 . It is well-known that the standard cross product can be obtained from quaternionic multiplication by (3.38), and that $\langle u \times v, w \rangle = \mu(u, v, w)$ is the volume form μ evaluated on the 3-plane $u \wedge v \wedge w$. Equivalently, the cross product is given by the Hodge star on \mathbb{R}^3 . That is, $u \times v = \star(u \wedge v)$. In this case $\star\varphi = \star\mu = 1$.
- (iii) The cross product on \mathbb{R}^7 is induced in the same way from octonionic multiplication, and is called the *exceptional cross product*. In this case φ is a nontrivial 3-form on \mathbb{R}^7 , and $\star\varphi = \psi$ is a nontrivial 4-form on \mathbb{R}^7 . We discuss these in more detail in Section 4.1. \blacktriangle

3.4 Further properties of the cross product in \mathbb{R}^3 and \mathbb{R}^7

Let us investigate some further properties of the cross product. First, note that for $\mathbb{V} = \mathbb{R}^3 \cong \text{Im } \mathbb{H}$, equation (3.51) reduces to the familiar $a \times (b \times c) = -\langle a, b \rangle c + \langle a, c \rangle b$, because \mathbb{H} is associative. However, for $\mathbb{V} = \mathbb{R}^7 \cong \text{Im } \mathbb{O}$, we have

$$a \times (b \times c) = -\langle a, b \rangle c + \langle a, c \rangle b - \frac{1}{2}[a, b, c] \quad (3.62)$$

where the last term *does not vanish in general*. In fact using (3.36) we can write (3.62) as

$$a \times (b \times c) = -\langle a, b \rangle c + \langle a, c \rangle b + (\psi(a, b, c, \cdot))^\sharp \quad (3.63)$$

where α^\sharp is the vector in \mathbb{V} that is metric-dual to the 1-form $\alpha \in \mathbb{V}^*$ via the inner product. Explicitly, $\langle \alpha^\sharp, b \rangle = \alpha(b)$ for all $b \in \mathbb{V}$.

Remark 3.64. The nontriviality of the last term in (3.62) or (3.63) is equivalent to the nonassociativity of \mathbb{O} and is the source of the inherent *nonlinearity* in geometries defined using the octonions. See also Remark 4.30 below. \blacktriangle

We obtain a number of important consequences from the fundamental identity (3.62). The remaining results in this section hold for both the cases $\mathbb{V} = \mathbb{R}^3 \cong \text{Im } \mathbb{H}$ and $\mathbb{V} = \mathbb{R}^7 \cong \text{Im } \mathbb{O}$, with the understanding that the associator term vanishes in the \mathbb{R}^3 case.

Corollary 3.65. *Let $a, c \in \mathbb{V}$. Then we have*

$$a \times (a \times c) = -\|a\|^2 c + \langle a, c \rangle a. \quad (3.66)$$

Proof. Let $a = b$ in (3.62). The associator term vanishes by Proposition 3.31. \square

Remark 3.67. From Corollary 3.65 we deduce the following observation. Let $a \in \mathbb{V}$ satisfy $\|a\| = 1$. Consider the codimension one subspace $\mathbb{U} = (\text{span}\{a\})^\perp$ orthogonal to a . Since $a \times c$ is orthogonal to c for all c , the linear map $J_a : \mathbb{V} \rightarrow \mathbb{V}$ given by $J_a(c) = a \times c$ leaves \mathbb{U} invariant, and by (3.66) we have $(J_a)^2 = -I$ on \mathbb{U} , so J_a is a complex structure on \mathbb{U} . \blacktriangle

Corollary 3.68. *Let $a, b, c \in \mathbb{V}$ be orthonormal, with $a \times b = c$. Then $b \times c = a$ and $c \times a = b$.*

Proof. Take the cross product of $a \times b = c$ on both sides with a or b and use (3.66). \square

Corollary 3.69. *Let $a, b, c, d \in \mathbb{V}$. Recall that*

$$\langle a \wedge b, c \wedge d \rangle = \det \begin{pmatrix} \langle a, c \rangle & \langle a, d \rangle \\ \langle b, c \rangle & \langle b, d \rangle \end{pmatrix} = \langle a, c \rangle \langle b, d \rangle - \langle a, d \rangle \langle b, c \rangle.$$

Then we have

$$\langle a \times b, c \times d \rangle = \langle a \wedge b, c \wedge d \rangle - \frac{1}{2} \langle a, [b, c, d] \rangle, \quad (3.70)$$

$$\langle a \times b, a \times d \rangle = \langle a \wedge b, a \wedge d \rangle = \|a\|^2 \langle b, d \rangle - \langle a, b \rangle \langle a, d \rangle. \quad (3.71)$$

Proof. Equation (3.71) follows from (3.70) by setting $c = a$ and using Proposition 3.33. To establish (3.70), we compute using (3.45) and the skew-symmetry of φ as follows:

$$\langle a \times b, c \times d \rangle = \varphi(a, b, c \times d) = -\varphi(a, c \times d, b) = -\langle a \times (c \times d), b \rangle.$$

Using (3.62), the above expression becomes

$$\begin{aligned} \langle a \times b, c \times d \rangle &= -(-\langle a, c \rangle d + \langle a, d \rangle c - \frac{1}{2} [a, c, d], b) \\ &= \langle a, c \rangle \langle b, d \rangle - \langle a, d \rangle \langle b, c \rangle + \frac{1}{2} \langle b, [a, c, d] \rangle. \end{aligned}$$

Using Proposition 3.33, the above expression equals (3.70). \square

Remark 3.72. Using (3.36), when $n = 7$ we can also write (3.70) as

$$\langle a \times b, c \times d \rangle = \langle a \wedge b, c \wedge d \rangle - \psi(a, b, c, d). \quad (3.73)$$

Recall from Remark 3.64 that the nontriviality of ψ is equivalent to the nonassociativity of \mathbb{O} . Thus the above equation says that the nonassociativity of \mathbb{O} is also equivalent to the fact that

$$\langle a \times b, c \times d \rangle \neq \langle a \wedge b, c \wedge d \rangle$$

in general.

By contrast, when $n = 3$ the associator vanishes, and we do have $\langle a \times b, c \times d \rangle = \langle a \wedge b, c \wedge d \rangle$ in this case. This corresponds, by (ii) of Remark 3.61, to the fact that $a \times b = \star(a \wedge b)$ and \star is an isometry. \blacktriangle

4 The geometry of G_2 -structures

In this section we discuss G_2 -structures, first on \mathbb{R}^7 and then on smooth 7-manifolds, including a discussion of the decomposition of the space of differential forms and of the torsion of a G_2 -structure.

4.1 The canonical G_2 -structure on \mathbb{R}^7

In this section we describe in more detail the canonical G_2 -structure on $\mathbb{R}^7 \cong \text{Im } \mathbb{O}$. This standard “ G_2 -package” on \mathbb{R}^7 consists of the standard Euclidean metric g_o , for which the standard basis e_1, \dots, e_7 is orthonormal, the standard volume form $\mu_o = e^1 \wedge \dots \wedge e^7$ associated to g_o and the standard orientation, the “associative” 3-form φ_o , the “coassociative” 4-form ψ_o , and finally the “cross product” \times_o operation. We use the “o” subscript for the standard G_2 -package $(g_o, \mu_o, \varphi_o, \psi_o, \times_o)$ on \mathbb{R}^7 to distinguish it from a general G_2 -structure on a smooth 7-manifold which is defined in Section 4.2. We also use $\|\cdot\|_o$ to denote both the norm on \mathbb{R}^7 induced from the inner product g_o and also the induced norm on $\Lambda^\bullet(\mathbb{R}^7)^*$.

We identify $\mathbb{R}^7 \cong \text{Im } \mathbb{O}$. Recall from Definition 3.34 that the associative 3-form φ_o and the coassociative 4-form ψ_o are given by

$$\begin{aligned}\varphi_o(a, b, c) &= \frac{1}{2}\langle [a, b], c \rangle && \text{for } a, b, c \in \mathbb{R}^7, \\ \psi_o(a, b, c, d) &= -\frac{1}{2}\langle [a, b, c], d \rangle && \text{for } a, b, c, d \in \mathbb{R}^7.\end{aligned}$$

Using the octonion multiplication table, one can show that with respect to the standard dual basis e^1, \dots, e^7 on $(\mathbb{R}^7)^*$, and writing $e^{ijk} = e^i \wedge e^j \wedge e^k$ and similarly for decomposable 4-forms, we have

$$\begin{aligned}\varphi_o &= e^{123} - e^{167} - e^{527} - e^{563} - e^{415} - e^{426} - e^{437}, \\ \psi_o &= e^{4567} - e^{4523} - e^{4163} - e^{4127} - e^{2637} - e^{1537} - e^{1526}.\end{aligned}\tag{4.1}$$

It is immediate that

$$\psi_o = *_o \varphi_o,$$

where $*_o$ is the Hodge star operator induced from (g_o, μ_o) . The explicit expressions for φ_o and $\psi_o = *_o \varphi_o$ in (4.1) are not enlightening and need not be memorized by the reader. There is a particular method to the seeming madness in which we have written φ_o and ψ_o , which is explained in [27] in relation to the standard $\text{SU}(3)$ -structure on $\mathbb{R}^7 = \mathbb{C}^3 \oplus \mathbb{R}$, where $z^1 = x^1 + ix^5$, $z^2 = x^2 + ix^6$, $z^3 = x^3 + ix^7$ are the complex coordinates on \mathbb{C}^3 and x^4 is the coordinate on \mathbb{R} .

One piece of information to retain from (4.1) is that

$$\|\varphi_o\|_o^2 = \|\psi_o\|_o^2 = 7,\tag{4.2}$$

which is equivalent to the identity $\varphi_o \wedge \psi_o = 7\mu_o$. (These facts are analogous to the identities $\|\omega_o\|_o^2 = 2m$ and $\frac{1}{m!}\omega_o^m = \mu_o$ for the standard Kähler form ω_o on \mathbb{C}^m with respect to the Euclidean metric.)

We now use this standard ‘‘G₂-package’’ on \mathbb{R}^7 to give a definition of the group G₂.

Definition 4.3. The group G₂ is defined to be the subgroup of $\text{GL}(7, \mathbb{R})$ that preserves the standard G₂-package on \mathbb{R}^7 . That is,

$$\text{G}_2 = \{A \in \text{GL}(7, \mathbb{R}) : A^* g_o = g_o, A^* \mu_o = \mu_o, A^* \varphi_o = \varphi_o\}.$$

Note that because g_o and μ_o determine the Hodge star operator $*_o$, which in turn from φ_o determines ψ_o , and because g_o and φ_o together determine \times_o , it follows that any $A \in \text{G}_2$ also preserves ψ_o and \times_o . (But see Theorem 4.4 below.) Moreover, since by definition $A \in \text{G}_2$ preserves the standard Euclidean metric and orientation on \mathbb{R}^7 , we see that G₂ as defined above is a *subgroup* of $\text{SO}(7, \mathbb{R})$. \blacktriangle

Theorem 4.4 (Bryant [4]). *Define $K = \{A \in \text{GL}(7, \mathbb{R}) : A^* \varphi_o = \varphi_o\}$. Then in fact $K = \text{G}_2$. That is, if $A \in \text{GL}(7, \mathbb{R})$ preserves φ_o , then it also automatically preserves g_o and μ_o as well.*

Proof. One can show using the explicit form (4.1) for φ_o in terms of the standard basis e^1, \dots, e^7 of $(\mathbb{R}^7)^*$ that

$$(a \lrcorner \varphi_o) \wedge (b \lrcorner \varphi_o) \wedge \varphi_o = -6g_o(a, b)\mu_o.\tag{4.5}$$

It follows from (4.5) that if $A^* \varphi_o = \varphi_o$, then

$$(A^* g_o)(a, b)A^* \mu_o = g_o(Aa, Ab)(\det A)\mu_o = g_o(a, b)\mu_o.\tag{4.6}$$

Thus we have $(\det A)g_o(Aa, Ab) = g_o(a, b)$, or equivalently in terms of matrices, $g_o = (\det A)A^T g_o A$. Taking determinants of both sides, and observing that these are all 7×7 matrices, gives $\det g_o = (\det A)^9 \det g_o$, so $\det A = 1$ and $A^* \mu_o = \mu_o$. But then (4.6) says that $A^* g_o = g_o$ as claimed. \square

Remark 4.7. In Bryant [4] the equation (4.5) has a +6 on the right hand side rather than our -6, because of a different orientation convention. See also Remark 4.37 below. \blacktriangle

Remark 4.8. Theorem 4.4 appears in [4]. Robert Bryant claims that it is a much older result, due to Élie Cartan. While this is almost certainly true, most mathematicians know this result as “Bryant’s Theorem” as [4] is the earliest accessible reference for this result that is widely known. See Agricola [1] for more about this history of the group G_2 . \blacktriangle

Corollary 4.9. *The group G_2 can equivalently be defined as the automorphism group $\text{Aut}(\mathbb{O})$ of the normed division algebra \mathbb{O} of octonions.*

Proof. Let $A \in \text{Aut}(\mathbb{O})$. Since A is an algebra automorphism we have $A(1) = 1$ and thus $A(t1) = t$ for all $t \in \mathbb{R}$. Now suppose $p \in \text{Im } \mathbb{O}$. Then $p^2 = -p\bar{p} = -\|p\|_{\mathbb{O}}^2$ is real. Thus we have

$$(A(p))^2 = A(p)A(p) = A(p^2) = A(-\|p\|_{\mathbb{O}}^2) = -\|p\|_{\mathbb{O}}^2$$

is real. By Corollary 3.25 we deduce that $A(p)$ must be real or imaginary. Suppose it is real. Then $A(p) = t1$ for some $t \in \mathbb{R}$. But then $A(p) = A(t1)$ and $p \neq t1$ since p is imaginary. This contradicts the invertibility of A . Thus $A(p)$ must be imaginary. This means $\overline{A(p)} = -A(p)$ whenever p is imaginary.

Now let $p = (\text{Re } p)1 + (\text{Im } p)$. Since A is linear over \mathbb{R} and $A(1) = 1$, we get $A(p) = (\text{Re } p)1 + A(\text{Im } p)$. But then $\overline{A(p)} = (\text{Re } p)1 - A(\text{Im } p) = A(\bar{p})$. It follows that

$$\|A(p)\|_{\mathbb{O}}^2 = A(p)\overline{A(p)} = A(p)A(\bar{p}) = A(p\bar{p}) = A(\|p\|_{\mathbb{O}}^2) = \|p\|_{\mathbb{O}}^2.$$

Thus $\|A(p)\|_{\mathbb{O}} = \|p\|_{\mathbb{O}}$, and from $A(1) = 1$ and $A(\text{Im } \mathbb{O}) \subseteq (\text{Im } \mathbb{O})$ we conclude that $A \in O(7)$. Finally, from (3.46), if $a, b, c \in \text{Im } \mathbb{O}$ we get

$$\begin{aligned} (A^* \varphi_{\mathbb{O}})(a, b, c) &= \varphi_{\mathbb{O}}(Aa, Ab, Ac) = \langle (Aa)(Ab), Ac \rangle \\ &= \langle A(ab), Ac \rangle = \langle ab, c \rangle = \varphi_{\mathbb{O}}(a, b, c). \end{aligned}$$

Thus $A^* \varphi_{\mathbb{O}} = \varphi_{\mathbb{O}}$, so by Theorem 4.4 we deduce that $A \in G_2$.

Conversely, if $A \in G_2$, then A preserves the cross product and the inner product, so if we extend A linearly from $\mathbb{R}^7 \cong \text{Im } \mathbb{O}$ to $\mathbb{O} = \mathbb{R} \oplus \mathbb{R}^7$ by setting $A(1) = 1$, then it follows immediately from (3.60) that $A(ab) = A(a)A(b)$ for all $a, b \in \mathbb{O}$, so $A \in \text{Aut}(\mathbb{O})$. \square

Remark 4.10. Theorem 4.4 is an *absolutely crucial* ingredient of G_2 -geometry. It says that the 3-form $\varphi_{\mathbb{O}}$ *determines* the orientation $\mu_{\mathbb{O}}$ and the metric $g_{\mathbb{O}}$ in a *highly nonlinear way*. This is in stark contrast to the situation of the standard $U(m)$ -structure on \mathbb{C}^m , which consists of the Euclidean metric $g_{\mathbb{O}}$, the standard complex structure $J_{\mathbb{O}}$ on \mathbb{C}^m , and the associated Kähler form $\omega_{\mathbb{O}}$, which are all related by

$$\omega_{\mathbb{O}}(a, b) = g_{\mathbb{O}}(J_{\mathbb{O}}a, b). \quad (4.11)$$

Moreover, the standard volume form is $\mu_{\mathbb{O}} = \frac{1}{m!} \omega_{\mathbb{O}}^m$. Equation (4.11) should be compared to (3.45). The almost complex structure $J_{\mathbb{O}}$ is the analogue of the cross product $\times_{\mathbb{O}}$, and the 2-form $\omega_{\mathbb{O}}$ is the analogue of the 3-form $\varphi_{\mathbb{O}}$. However, in the case of the standard $U(m)$ -structure, the 2-form $\omega_{\mathbb{O}}$ *does not* determine the metric. (Although it *does* determine the orientation.) The correct way to think about (4.11) is that knowledge of any two of $g_{\mathbb{O}}, J_{\mathbb{O}}, \omega_{\mathbb{O}}$ uniquely determines the third. This is encoded by the following Lie group relation:

$$O(2m, \mathbb{R}) \cap \text{GL}(m, \mathbb{C}) = O(2m, \mathbb{R}) \cap \text{Sp}(m, \mathbb{R}) = \text{GL}(m, \mathbb{C}) \cap \text{Sp}(m, \mathbb{R}) = U(m).$$

Colloquially, we say that the intersection of any two of Riemannian, complex, and symplectic geometry is Kähler geometry. By contrast, G_2 geometry *does not “decouple”* in any such way. It is *not* the intersection of Riemannian geometry with any other “independent” geometry. The 3-form $\varphi_{\mathbb{O}}$ determines *everything else*. \blacktriangle

Let us consider how we should think about the group G_2 , which by Theorem 4.4 is described as a particular *subgroup* of $\text{SO}(7, \mathbb{R})$. Before we can do that, we need a preliminary result.

Lemma 4.12. *Let f_1, f_2, f_4 be a triple of orthonormal vectors in \mathbb{R}^7 such that $\varphi_o(f_1, f_2, f_4) = 0$. Define*

$$f_3 = f_1 \times_o f_2, \quad f_5 = f_1 \times_o f_4, \quad f_6 = f_2 \times_o f_4, \quad f_7 = f_3 \times_o f_4 = (f_1 \times_o f_2) \times_o f_4. \quad (4.13)$$

Then the ordered set $\{f_1, \dots, f_7\}$ is an oriented orthonormal basis of \mathbb{R}^7 .

Proof. One can check using equations (3.41), (3.45), and (3.71), together with the hypotheses that $\{f_1, f_2, f_4\}$ are orthonormal and $\varphi_o(f_1, f_2, f_4) = 0$, that $\langle f_i, f_j \rangle = \delta_{ij}$ for all i, j so the set is orthonormal. Most of these are immediate. We demonstrate one of the less trivial cases. Using Corollary 3.68, we deduce that $f_3 \times_o f_1 = f_2$. Thus we have

$$\begin{aligned} g_o(f_1, f_7) &= g_o(f_1, f_3 \times_o f_4) = \varphi_o(f_1, f_3, f_4) \\ &= -\varphi_o(f_3, f_1, f_4) = -g_o(f_3 \times_o f_1, f_4) = -g_o(f_2, f_4) = 0. \end{aligned}$$

It remains to show $\{f_1, \dots, f_7\}$ induces the same orientation as $\{e_1, \dots, e_7\}$. When $f_k = e_k$ for $k = 1, 2, 4$, then it follows from the octonion multiplication table and (4.13) that $f_k = e_k$ for all $k = 1, \dots, 7$. It is then not hard to see that the matrix in $A \in O(7)$ given by

$$A = (f_1 | f_2 | f_3 | f_4 | f_5 | f_6 | f_7)$$

can be obtained from the identity matrix by a product of three elements of $SO(7)$. Thus $A \in SO(7)$ and hence $\{f_1, \dots, f_7\}$ is oriented. \square

Corollary 4.14. *The group G_2 can be viewed explicitly as the subgroup of $SO(7)$ consisting of those elements $A \in SO(7)$ of the form*

$$A = (f_1 | f_2 | f_1 \times_o f_3 | f_4 | f_1 \times_o f_4 | f_2 \times_o f_4 | (f_1 \times_o f_2) \times_o f_4) \quad (4.15)$$

where $\{f_1, f_2, f_4\}$ is an orthonormal triple satisfying $\varphi_o(f_1, f_2, f_4) = 0$. (This means that the cross product of any two of $\{f_1, f_2, f_4\}$ is orthogonal to the third.)

Proof. By Lemma 4.12, every matrix of the form (4.15) does lie in $SO(7)$. By Theorem 4.4, a matrix $A \in SO(7)$ is in G_2 if and only if A preserves the vector cross product \times_o . Since A takes e_k to f_k , it follows from the fact that

$$e_3 = e_1 \times_o e_2, \quad e_5 = e_1 \times_o e_4, \quad e_6 = e_2 \times_o e_4, \quad e_7 = e_3 \times_o e_4 = (e_1 \times_o e_2) \times_o e_4,$$

that such the elements of G_2 are precisely the matrices of the form (4.15). \square

Remark 4.16. We can argue from Corollary 4.14 that $\dim G_2 = 14$, as follows. We know G_2 corresponds to the set of orthonormal triples $\{f_1, f_2, f_4\}$ such that f_4 is orthogonal to f_1, f_2 , and $f_1 \times_o f_2$. Thus f_1 is any unit vector in \mathbb{R}^7 , so it lies on S^6 . Then f_2 must be orthogonal to f_1 , so it lies on the unit sphere S^5 of the \mathbb{R}^6 that is orthogonal to f_1 . Finally, f_4 must be orthogonal to f_1, f_2 , and $f_1 \times_o f_2$, so it lies on the unit sphere S^3 of the \mathbb{R}^4 that is orthogonal to these three vectors. Thus we have $6 + 5 + 3 = 14$ degrees of freedom, so $\dim G_2 = 14$.

(In fact, G_2 is a connected, simply-connected, compact Lie subgroup of $SO(7)$.) \blacktriangle

4.2 G_2 -structures on smooth 7-manifolds

In this section, as discussed in Section 2, we equip a smooth 7-manifold with the “ G_2 package” at each tangent space, in a smoothly varying way.

Definition 4.17. Let M^7 be a smooth 7-manifold. A G_2 -structure on M is a smooth 3-form φ on M such that, at every $p \in M$, there exists a linear isomorphism $T_p M \cong \mathbb{R}^7$ with respect to which $\varphi_p \in \Lambda^3(T_p^* M)$ corresponds to $\varphi_o \in \Lambda^3(\mathbb{R}^7)^*$. Therefore, because φ_o induces g_o and μ_o , a G_2 -structure φ on M induces a Riemannian metric g_φ and associated Riemannian volume form μ_φ . These in turn induce a Hodge star operator \star_φ and dual 4-form $\psi = \star_\varphi \varphi$. \blacktriangle

Thus if φ is a G_2 -structure on M , then at every point $p \in M$, there exists a basis $\{e_1, \dots, e_7\}$ of $T_p M$ with respect to which $\varphi_p = \varphi_o$ from (4.1). Note that in general we *cannot* choose a *local frame* on an open set U in M with respect to which φ takes the standard form in (4.1), *we can only do this at a single point*. This is analogous to how, in a manifold (M^{2m}, g, J, ω) with $U(m)$ -structure, we can always find a basis of $T_p M$ for any $p \in M$ in which the “ $U(m)$ package” assumes the standard form on \mathbb{C}^m , but we cannot in general do this on an open set. (See [12] for a comprehensive treatment of $U(m)$ -structures.)

Not every smooth 7-manifold admits G_2 -structures. A G_2 -structure is equivalent to a reduction of the structure group of the frame bundle of M from $GL(7, \mathbb{R})$ to $G_2 \subset SO(7)$. As such, the existence of a G_2 -structure is entirely a topological question.

Proposition 4.18. *A smooth 7-manifold M admits a G_2 -structure if and only if M is both orientable and spinnable. This is equivalent to the vanishing of the first two Stiefel-Whitney classes $w_1(TM)$ and $w_2(TM)$.*

Proof. See Lawson–Michelsohn [31, Chapter IV, Theorem 10.6] for a proof. \square

Therefore, while not all smooth 7-manifolds admit G_2 -structures, there are many that do and they are completely characterized by Proposition 4.18.

There is a much more concrete way to understand when a 3-form φ on M is a G_2 -structure. It can be considered as a “working differential geometer’s definition of G_2 -structure”, and is described as follows. Let $\varphi \in \Omega^3(M^7)$. Let x^1, \dots, x^7 be local coordinates on an open set U in M . For $i, j \in \{1, \dots, 7\}$, define a smooth function B_{ij} on U by

$$-6B_{ij} dx^1 \wedge \dots \wedge dx^7 = \left(\frac{\partial}{\partial x^i} \lrcorner \varphi \right) \wedge \left(\frac{\partial}{\partial x^j} \lrcorner \varphi \right) \wedge \varphi. \quad (4.19)$$

Since 2-forms commute, we have $B_{ij} = B_{ji}$. In fact, comparison with (4.5) shows that if φ is a G_2 -structure, we must have $B_{ij} = g_{ij} \sqrt{\det g}$, since $\mu = \sqrt{\det g} dx^1 \wedge \dots \wedge dx^7$ is the Riemannian volume form in local coordinates. Hence $\det B = (\sqrt{\det g})^7 \det g = (\det g)^{\frac{9}{2}}$ and thus $\sqrt{\det g} = (\det B)^{\frac{1}{9}}$. Solving for g_{ij} gives

$$g_{ij} = \frac{1}{(\det B)^{\frac{1}{9}}} B_{ij}. \quad (4.20)$$

We say that $\varphi \in \Omega^3(M^7)$ is a G_2 -structure if this recipe *actually works* to construct a Riemannian metric. Thus we must have both:

- (i) $\det B$ must be nonzero everywhere on U ,
- (ii) g_{ij} as defined in (4.20) must be positive definite everywhere on U .

Of course, these two conditions must hold in any local coordinates x^1, \dots, x^7 on M . But the advantage of this way of thinking about G_2 -structures (besides it being very concrete) is that it allows us to easily see that the condition of φ being a G_2 -structure is an *open condition*. That is, if φ is a G_2 -structure, and $\tilde{\varphi}$ is another smooth 3-form on M sufficiently close to φ (in the C^0 -norm with respect to any Riemannian metric on M) then $\tilde{\varphi}$ will also be a G_2 -structure. This is because both conditions (i) and (ii) above are open conditions at each point p of M .

We conclude that, if the space of G_2 -structures on M is *nonempty*, then it can be identified with a space Ω_+^3 of smooth sections of a fibre bundle $\Lambda_+^3(T^*M)$ whose fibres are open subsets of the corresponding

fibres of the bundle $\Lambda^3(T^*M)$. The space Ω_+^3 is also called the space of *nondegenerate* or *positive* or *stable* 3-forms on M .

Remark 4.21. There is another way of seeing that the condition of being a G_2 -structure is open. At any point $p \in M$, the space of all G_2 -structures $\Lambda_+^3(T_p^*M)$ can be identified with the orbit of φ_o in $\Lambda^3(\mathbb{R}^7)^*$ by the action of $\text{GL}(7, \mathbb{R})$ quotiented by the stabilizer subgroup of φ_o , which is G_2 by Theorem 4.4. Since $\dim \text{GL}(7, \mathbb{R}) = 49$, and $\dim G_2 = 14$, we have $\dim \Lambda_+^3(T_p^*M) = 49 - 14 = 35 = \dim \Lambda^3(T_p^*M)$, and thus $\Lambda_+^3(T_p^*M)$ is an open set of $\Lambda^3(T_p^*M)$. See Hitchin [20] for a general discussion of *stable forms*. \blacktriangle

Remark 4.22. The nonlinear map $\varphi \rightarrow g$ is not one-to-one. In fact, given a metric g on M induced from a G_2 -structure φ , at each point $p \in M$, the space of G_2 -structures at p inducing g_p is diffeomorphic to \mathbb{RP}^7 . Thus the G_2 -structures inducing the same metric g correspond to sections of an \mathbb{RP}^7 -bundle over M . See [6, page 10, Remark 4] for more details on *isometric G_2 -structures*. \blacktriangle

Let (M, φ) be a manifold with G_2 -structure, and let g be the induced metric. Let $\psi = \star_\varphi \varphi$ denote the dual 4-form. The vital relation (3.63), which is equivalent to (3.73) leads to fundamental local coordinate identities relating φ , ψ , and g .

Theorem 4.23. *In local coordinates on M , the tensors φ , ψ , and g satisfy the following relations:*

$$\varphi_{ijk}\varphi_{abc}g^{kc} = g_{ia}g_{jb} - g_{ib}g_{ja} - \psi_{ijab}, \quad (4.24)$$

$$\varphi_{ijk}\varphi_{abc}g^{jb}g^{kc} = 6g_{ia}, \quad (4.25)$$

$$\varphi_{ijk}\psi_{abcd}g^{kd} = g_{ia}\varphi_{jbc} + g_{ib}\varphi_{ajc} + g_{ic}\varphi_{abj} - g_{aj}\varphi_{ibc} - g_{bj}\varphi_{aic} - g_{cj}\varphi_{abi}, \quad (4.26)$$

$$\varphi_{ijk}\psi_{abcd}g^{jc}g^{kd} = -4\varphi_{iab}, \quad (4.27)$$

$$\psi_{ijkl}\psi_{abcd}g^{kc}g^{ld} = 4g_{ia}g_{jb} - 4g_{ib}g_{ja} - 2\psi_{ijab}, \quad (4.28)$$

$$\psi_{ijkl}\psi_{abcd}g^{jb}g^{kc}g^{ld} = 24g_{ia}. \quad (4.29)$$

Proof. These are derived from the relation (3.63) or equivalently (3.73). Indeed, the first identity (4.24) is precisely (3.73) expressed in local coordinates. The explicit details can be found in [26, Section A.3]. \square

Of course, there are many other possible contractions of φ , ψ , and g . In Theorem 4.23 we only list those that show up most frequently in practice.

Remark 4.30. The identities for G_2 -structures in Theorem 4.23 should be contrasted with the analogue for $U(m)$ -structures. First, we have only a single form ω , as opposed to the two forms φ and ψ . Moreover, from $\omega_{ab} = J_a^c g_{cb}$, which comes from (4.11), and the fact that $J^2 = -I$, we find that $\omega_{ia}\omega_{jb}g^{ab} = g_{ij}$. This is much simpler than (4.24) as the right hand side only involves the metric g . This again illustrates the “increased nonlinearity” of G_2 geometry, as mentioned in Remark 3.64 above. \blacktriangle

4.3 Decomposition of Ω^\bullet into irreducible G_2 representations

Let (M, φ) be a manifold with G_2 -structure. The bundle $\Lambda^\bullet(T^*M) = \oplus_{k=1}^7 \Lambda^k(T^*M)$ decomposes into irreducible representations of G_2 . This in turn induces a decomposition of the space $\Omega^k = \Gamma(\Lambda^k(T^*M))$ of smooth k -forms on M . This is entirely analogous to how, on a manifold with almost complex structure, the space $\Omega_{\mathbb{C}}^\bullet = \Gamma(\Lambda^\bullet(T^*M) \otimes \mathbb{C})$ of complex-valued forms decomposes into “forms of type (p, q) ”.

By Theorem 4.4, all the tensors determined by φ will be invariant under G_2 and hence any subspaces of Ω^k defined using φ , ψ , g , and \star will be G_2 representations. The space Ω^k is *irreducible* if $k = 0, 1, 6, 7$. However, for $k = 2, 3, 4, 5$ we have a nontrivial decomposition. Since $\Omega^k = \star \Omega^{7-k}$, the decompositions of Ω^5 and Ω^4 are obtained by taking \star of the decompositions of Ω^2 and Ω^3 , respectively.

In fact we have

$$\begin{aligned}\Omega^2 &= \Omega_7^2 \oplus \Omega_{14}^2, \\ \Omega^3 &= \Omega_1^3 \oplus \Omega_7^3 \oplus \Omega_{27}^3,\end{aligned}$$

where Ω_l^k has (pointwise) dimension l and these decompositions are *orthogonal* with respect to g . These spaces are described invariantly as follows:

$$\begin{aligned}\Omega_7^2 &= \{X \lrcorner \varphi \mid X \in \Gamma(TM)\} = \{\beta \in \Omega^2 \mid \star(\varphi \wedge \beta) = -2\beta\}, \\ \Omega_{14}^2 &= \{\beta \in \Omega^2 \mid \beta \wedge \psi = 0\} = \{\beta \in \Omega^2 \mid \star(\varphi \wedge \beta) = \beta\},\end{aligned}\tag{4.31}$$

and

$$\begin{aligned}\Omega_1^3 &= \{f\varphi \mid f \in \Omega^0\}, & \Omega_7^3 &= \{X \lrcorner \psi \mid X \in \Gamma(TM)\}, \\ \Omega_{27}^3 &= \{\gamma \in \Omega^3 \mid \gamma \wedge \varphi = 0, \gamma \wedge \psi = 0\}.\end{aligned}\tag{4.32}$$

It is sometimes necessary to get our hands dirty, so we need to describe these subspaces in terms of local coordinates. Consider first the G_2 -invariant linear map $P : \Omega^2 \rightarrow \Omega^2$ given by $P\beta = \star(\varphi \wedge \beta)$. If we write $\beta = \frac{1}{2}\beta_{ij}dx^i \wedge dx^j$ and $P\beta = \frac{1}{2}(P\beta)_{ab}dx^a dx^b$, then one can show [26, Section 2.2] that

$$(P\beta)_{ab} = \frac{1}{2}\psi_{abcd}g^{ci}g^{dj}\beta_{ij}.\tag{4.33}$$

That is, up to the factor of $\frac{1}{2}$, the map P is given by contracting the 2-form with the 4-form ψ on two indices. It is easy to check that P is self-adjoint and thus orthogonally diagonalizable with real eigenvalues. Using the fundamental identity (4.28) for the contraction of ψ with itself on two indices, we find

$$\begin{aligned}(P^2\beta)_{ab} &= \frac{1}{2}\psi_{abcd}g^{ci}g^{dj}(P\beta)_{ij} = \frac{1}{4}\psi_{abcd}g^{ci}g^{dj}\psi_{ijst}g^{sp}g^{tq}\beta_{pq} \\ &= \frac{1}{4}(4g_{as}g_{bt} - 4g_{at}g_{bs} - 2\psi_{abst})g^{sp}g^{tq}\beta_{pq} \\ &= \beta_{ab} - \beta_{ba} - \frac{1}{2}\psi_{abst}g^{sp}g^{tq}\beta_{pq} = 2\beta_{ab} - (P\beta)_{ab}.\end{aligned}$$

Thus we deduce that $P^2 = 2I - P$, so $(P + 2I)(P - I) = 0$. Thus the eigenvalues of P are -2 and $+1$, in agreement with (4.31). To verify that $\lambda = -2$ corresponds to Ω_7^2 as given in (4.31), we let $\beta_{ij} = (X \lrcorner \varphi)_{ij} = X^m \varphi_{mij}$. Then using (4.27) we have

$$(P\beta)_{ab} = \frac{1}{2}\psi_{abcd}g^{ci}g^{dj}X^m \varphi_{mij} = -2X^m \varphi_{mab} = -2\beta_{ab},$$

as claimed. Also, the condition that $\Omega_{14}^2 = (\Omega_7^2)^\perp$ gives that $\beta \in \Omega_{14}^2$ must satisfy $X^m \varphi_{mij} \beta_{ab} g^{ia} g^{jb} = 0$ for all X^m . This is equivalent to $\varphi_{mij} \beta_{ab} g^{ia} g^{jb} = 0$. Thus, we can describe the decomposition (4.31) of Ω^2 in local coordinates as

$$\begin{aligned}\beta_{ij} \in \Omega_7^2 &\iff \beta_{ij} = X^m \varphi_{mij} &\iff \frac{1}{2}\psi_{abcd}g^{ci}g^{dj}\beta_{ij} = -2\beta_{ab}, \\ \beta_{ij} \in \Omega_{14}^2 &\iff \beta_{ij} g^{ia} g^{jb} \varphi_{abc} = 0 &\iff \frac{1}{2}\psi_{abcd}g^{ci}g^{dj}\beta_{ij} = \beta_{ab}.\end{aligned}\tag{4.34}$$

Moreover, it is easy to check using (4.25) that for $\beta \in \Omega_7^2$ we have

$$\beta_{ij} = X^m \varphi_{mij} \iff X^m = \frac{1}{6}\beta_{ab} g^{ai} g^{bj} \varphi_{ijk} g^{km}.\tag{4.35}$$

Remark 4.36. The description of the orthogonal splitting $\Omega^2 = \Omega_7^2 \oplus \Omega_{14}^2$ in terms of the $-2, +1$ eigenspaces of the operator $\beta \mapsto \star(\varphi \wedge \beta)$ is analogous to the orthogonal splitting $\Omega^2 = \Omega_+^2 \oplus \Omega_-^2$ into self-dual and anti-self-dual 2-forms on an oriented Riemannian 4-manifold with respect to the operator $\beta \mapsto \star\beta$. This analogy is important in G_2 gauge theory. See the lectures by Walpuski [41] in the present volume for an introduction to G_2 gauge theory. \blacktriangle

Remark 4.37. Many authors prefer to use the opposite orientation than we do for the orientation induced by φ . (See [27] for more details.) This changes the sign of \star . The upshot is that the eigenvalues $(-2, +1)$ in (4.31) and (4.34) are replaced by $(+2, -1)$. Readers should take care to be aware of any particular paper's sign conventions. \blacktriangle

The local coordinate description of the decomposition (4.32) of Ω^3 can be understood by considering the infinitesimal action of the $(1, 1)$ tensors $\Gamma(T^*M \otimes TM)$ on φ . Let $A = A_i^j \in \Gamma(T^*M \otimes TM)$. At each point $p \in M$, we have $e^{At} \in \text{GL}(T_pM)$, and thus

$$e^{At} \cdot \varphi = \frac{1}{6} \varphi_{ijk} (e^{At} dx^i) \wedge (e^{At} dx^j) \wedge (e^{At} dx^k). \quad (4.38)$$

Define $A \diamond \varphi \in \Omega^3$ by

$$(A \diamond \varphi) = \left. \frac{d}{dt} \right|_{t=0} (e^{At} \cdot \varphi). \quad (4.39)$$

From (4.38) we compute

$$(A \diamond \varphi) = \frac{1}{6} (A_i^l \varphi_{ljk} + A_j^l \varphi_{ilk} + A_k^l \varphi_{ijl}) dx^i \wedge dx^j \wedge dx^k,$$

and hence

$$(A \diamond \varphi)_{ijk} = A_i^l \varphi_{ljk} + A_j^l \varphi_{ilk} + A_k^l \varphi_{ijl}. \quad (4.40)$$

Use the metric g to identify $A \in \Gamma(T^*M \otimes TM)$ with a bilinear form $A \in \Gamma(T^*M \otimes T^*M)$ by $A_{ij} = A_i^l g_{lj}$. Recall from Section 1.2 that there is an orthogonal splitting

$$\Gamma(T^*M \otimes T^*M) \cong \Omega^0 \oplus \mathcal{S}_0 \oplus \Omega^2.$$

By the orthogonal decomposition (4.31) on Ω^2 discussed above, we can further decompose this as

$$\Gamma(T^*M \otimes T^*M) \cong \Omega^0 \oplus \mathcal{S}_0 \oplus \Omega_7^2 \oplus \Omega_{14}^2.$$

With respect to this splitting, we can write $A = \frac{1}{7}(\text{Tr } A)g + A_0 + A_7 + A_{14}$, where A_0 is a traceless symmetric tensor.

By (4.40), we have a linear map $A \mapsto A \diamond \varphi$ from $\Omega^0 \oplus \mathcal{S}_0 \oplus \Omega_7^2 \oplus \Omega_{14}^2$ to the space Ω^3 .

Proposition 4.41. *The kernel of $A \mapsto A \diamond \varphi$ is Ω_{14}^2 , and the remaining three summands Ω^0 , \mathcal{S}_0 , Ω_7^2 , of $\Gamma(T^*M \otimes T^*M)$ are mapped isomorphically onto Ω_1^3 , Ω_{27}^3 , Ω_7^3 , respectively. Explicitly, if $A = \frac{1}{7}(\text{Tr } A)g + A_0 + A_7 + A_{14}$, then*

$$A \diamond \varphi = \underbrace{\frac{3}{7}(\text{Tr } A)\varphi}_{\Omega_1^3} + \underbrace{A_0 \diamond \varphi}_{\Omega_{27}^3} + \underbrace{X \lrcorner \psi}_{\Omega_7^3},$$

where

$$X^m = -\frac{1}{2} A_{ij} g^{ia} g^{jb} \varphi_{abc} g^{cm}.$$

Proof. This can be established using the various contraction identities of Theorem 4.23. The explicit details can be found in [26, Section 2.2]. \square

Remark 4.42. The fact that Ω_{14}^2 is the kernel of $A \mapsto A \diamond \varphi$ is a consequence of the fact that G_2 is the Lie group that preserves φ . Thus the infinitesimal action, which is the action of the Lie algebra \mathfrak{g}_2 , annihilates φ . This is consistent with the fact that $G_2 \subset \text{SO}(7)$, so $\mathfrak{g}_2 \subset \mathfrak{so}(7) \cong \Lambda^2(\mathbb{R}^7)^*$. Thus, at every point $p \in M$, the space $\Lambda_{14}^2(T_p^*M)$ is isomorphic to \mathfrak{g}_2 . \blacktriangle

4.4 The torsion of a G_2 -structure

Let (M, φ) be a manifold with G_2 -structure. Since φ determines a Riemannian metric g , we get a Levi-Civita covariant derivative ∇ . Thus it makes sense to consider the tensor $\nabla\varphi \in \Gamma(T^*M \otimes \Lambda^3 T^*M)$.

Definition 4.43. The G_2 -structure φ is called *torsion-free* if $\nabla\varphi = 0$. Although this appears to be a linear equation, recall that ∇ is induced from g which itself depends nonlinearly on φ . Thus the equation $\nabla\varphi = 0$ is in fact a fully nonlinear first order partial differential equation for φ . We say (M, φ) is a *torsion-free G_2 manifold* if φ is a torsion-free G_2 -structure on M . For brevity, we sometimes use the term “ G_2 manifold” when we mean “torsion-free G_2 manifold”. \blacktriangle

The fundamental observation about the torsion of any G_2 -structure is the following.

Theorem 4.44. *Let X be a vector field on M . Then the 3-form $\nabla_X\varphi$ lies in the subspace Ω_7^3 of Ω^3 . Thus the covariant derivative $\nabla\varphi$ is a smooth section of $T^*M \otimes \Lambda_7^3(T^*M)$.*

Proof. By Proposition 4.41, any 3-form γ can be written as $\gamma = A \diamond \varphi$ for a unique $A = \frac{1}{7}(\text{Tr } A)g + A_0 + A_7$. We take the inner product of $A \diamond \varphi$ with $\nabla_X\varphi$. Using (4.40), this is

$$\begin{aligned} \langle A \diamond \varphi, \nabla_X\varphi \rangle &= \frac{1}{6}(A \diamond \varphi)_{ijk}(\nabla_X\varphi)_{abc}g^{ia}g^{jb}g^{kc} \\ &= \frac{1}{6}(A_i^l\varphi_{ljk} + A_j^l\varphi_{ilk} + A_k^l\varphi_{ijl})X^m\nabla_m\varphi_{abc}g^{ia}g^{jb}g^{kc} \\ &= \frac{1}{2}A_i^l\varphi_{ljk}X^m\nabla_m\varphi_{abc}g^{ia}g^{jb}g^{kc} \\ &= \frac{1}{2}A_{ip}X^m\varphi_{qjk}\nabla_m\varphi_{abc}g^{pq}g^{ia}g^{jb}g^{kc}. \end{aligned}$$

Taking the covariant derivative of (4.25) and using that g is parallel, we get

$$\nabla_m\varphi_{qjk}\varphi_{abc}g^{jb}g^{kc} = -\varphi_{qjk}\nabla_m\varphi_{abc}g^{jb}g^{kc}.$$

This says that $\nabla_m\varphi_{qjk}\varphi_{abc}g^{jb}g^{kc}$ is skew in q, a . Thus the symmetric part of A_{ip} does not contribute to $\langle A \diamond \varphi, \nabla_X\varphi \rangle$ above. That is, $\nabla_X\varphi$ is orthogonal to any element of $\Omega_1^3 \oplus \Omega_{27}^3$, as claimed. \square

Theorem 4.44 motivates the following definition.

Definition 4.45. Because $\nabla_X\varphi \in \Omega_7^3$, by (4.32) we can write

$$\nabla_X\varphi = T(X) \lrcorner \psi$$

for some vector field $T(X)$ on M . That is, there exists a tensor $T \in \Gamma(T^*M \otimes T^*M)$ such that

$$\nabla_m\varphi_{ijk} = T_{mp}g^{pq}\psi_{qijk}. \quad (4.46)$$

We call T the *full torsion tensor* of φ . \blacktriangle

By contracting (4.46) with ψ_{nijk} on i, j, k and using (4.29), we obtain

$$T_{mn} = \frac{1}{24}\nabla_m\varphi_{ijk}\psi_{nabc}g^{ia}g^{jb}g^{kc}. \quad (4.47)$$

Moreover, taking the covariant derivative of (4.24) and using (4.46) and (4.26), one can compute that

$$\nabla_p\psi_{ijkl} = -T_{pi}\varphi_{jkl} + T_{pj}\varphi_{ikl} - T_{pk}\varphi_{ijl} + T_{pl}\varphi_{ijk}. \quad (4.48)$$

Observe that equations (4.46) and (4.47) show that $\nabla\varphi = 0$ if and only if $T = 0$. (In this case (4.48) shows that $\nabla\psi = 0$ as well, which is also clear because $\psi = \star\varphi$ and ∇ commutes with \star .)

Hence φ is torsion-free if and only if $T = 0$. The tensor T is a more convenient measure of the failure of φ to be parallel, because we can easily decompose it into four independent pieces in $\Gamma(T^*M \otimes T^*M) \cong \Omega^0 \oplus \mathcal{S}_0 \oplus \Omega_7^2 \oplus \Omega_{14}^2$, as

$$T = T_1 + T_{27} + T_7 + T_{14}, \quad (4.49)$$

where $T_1 = \frac{1}{7}(\text{Tr } T)g$ and $T_{27} = T_0$ is the traceless symmetric part of T .

Corollary 4.50. *Let φ be a G_2 -structure on M . Then φ is torsion-free if and only if both $d\varphi = 0$ and $d\psi = 0$.*

Proof. Note that $d\psi = d \star \varphi = - \star d^* \varphi$, so $d\psi = 0$ if and only if $d^* \varphi = 0$. Because both the exterior derivative d and its adjoint d^* can be written in terms of ∇ , any parallel form is always closed and coclosed. It is the converse that is nontrivial here. In fact, $d\varphi$ and $d^* \varphi$ are *linear* in $\nabla \varphi$ and hence linear in T . Since $d\varphi \in \Omega^4 = \Omega_1^4 \oplus \Omega_7^4 \oplus \Omega_{27}^4$ and $d^* \varphi \in \Omega^2 = \Omega_7^2 \oplus \Omega_{14}^2$, it follows by Schur's Lemma that the independent components of $d\varphi$ and $d^* \varphi$ must correspond to the 1, 7, 14, 27 components of T as in (4.49), up to constant factors. Thus if $d\varphi = 0$ and $d^* \varphi = 0$, we must have $T = 0$. \square

Corollary 4.50 is a classical theorem of Fernández–Gray [14]. The present proof is an extremely abridged version of the argument in [26, Section 2.3].

Remark 4.51. Recall that a differential form γ on (M, g) is *harmonic* if $\Delta_d \gamma = (dd^* + d^*d)\gamma = 0$. On a compact manifold, by integration by parts harmonicity is equivalent to $d\gamma = 0$ and $d^* \gamma = 0$. Thus Corollary 4.50 says that in the compact case, a G_2 -structure φ is torsion-free if and only if it is harmonic with respect to its induced metric. \blacktriangle

Since the torsion T of φ decomposes into four independent components as in (4.49), each component can be zero or nonzero. This gives $2^4 = 16$ distinct classes of G_2 -structures. Some of the more interesting classes of G_2 -structures are given in the following table.

T_1	T_{27}	T_7	T_{14}	G_2 -structure
0	0	0	0	$\nabla \varphi = 0$ (torsion-free)
0	0	0	*	$d\varphi = 0$ (closed)
*	*	0	0	$d^* \varphi = 0$ (coclosed)
*	0	0	0	$d\varphi = \lambda \psi$ ($\lambda \neq 0$)

The last class in the table above is called *nearly parallel*, and one can show that λ is constant and that the induced metric is *positive Einstein*, with $R_{ij} = \frac{3}{8} \lambda^2 g_{ij}$. (For example, see [26, after Remark 4.19].)

More details on the 16 classes of G_2 -structures can be found in [8, 14, 25, 26]. In particular it is worth remarking [26, Theorem 2.32] that with respect to conformal changes of G_2 -structure, the component T_7 plays a very different role than the other three components T_1, T_{27}, T_{14} .

Aside. There is an equivalent approach to studying G_2 -structures using *spin geometry*. Let (M^7, g) be a Riemannian 7-manifold equipped with a spin structure and associated *spinor bundle* $\mathcal{S}(M)$. This is a real rank 8 vector bundle over M . Since $8 > 7$, by algebraic topology, this bundle always admits nowhere vanishing sections. Such a section s determines a 3-form φ on M by $\varphi(a, b, c) = \langle a \cdot b \cdot c \cdot s, s \rangle$, where \cdot denotes the Clifford multiplication of tangent vectors to M on spinors. Using the fact that s is nowhere zero, one can show that the 3-form φ is always a G_2 -structure. Moreover, φ is torsion-free if and only if s is a *parallel spinor*, with respect to the spin connection on $\mathcal{S}(M)$ induced from the Levi-Civita connection of g . (The existence of a parallel spinor for torsion-free G_2 manifolds is precisely why they are of interest in theoretical physics, as this is related to *supersymmetry*.) Similarly, φ is nearly parallel in the sense defined above if and only if s is a *Killing spinor*. The reader is directed to Harvey [17], Lawson–Michelsohn [31, Chapter IV. 10], and the more recent paper by Agricola–Chiossi–Friedrich–Höll [2] for more on this point of view. This approach is also very important in the construction of *invariants* of G_2 -structures, as discussed by Crowley–Goette–Nordström [11] in the present volume.

4.5 Relation between curvature and torsion for a G_2 -structure

Let (M, φ) be a manifold with G_2 -structure. Since φ determines a Riemannian metric g_φ , we have a Riemann curvature tensor R . There is an important relation between the tensors R and ∇T , called the “ G_2 Bianchi identity” that originally appeared in [26, Theorem 4.2].

Theorem 4.52. *The G_2 -Bianchi identity is the following:*

$$\nabla_i T_{jk} - \nabla_j T_{ik} = (T_{ip} T_{jq} + \frac{1}{2} R_{ijpq}) g^{pa} g^{qb} \varphi_{abk}. \quad (4.53)$$

Proof. Equation (4.53) can be derived by combining the covariant derivative of (4.46) with (4.48) to get an expression for $\nabla_m \nabla_p \varphi_{ijk}$ in terms of φ , ψ , and T , and ∇T . Then applying the Ricci identity to the difference

$$\nabla_m \nabla_p \varphi_{ijk} - \nabla_p \nabla_m \varphi_{ijk}$$

introduces Riemann curvature terms. Simplifying further using the identities of Theorem 4.23 eventually results in (4.53). \square

An important consequence of Theorem 4.52 is the following.

Corollary 4.54. *The Ricci curvature R_{jk} of the metric g induced by a G_2 -structure φ can be expressed in terms of the torsion T and its covariant derivative ∇T as follows:*

$$\begin{aligned} R_{jk} = & (\nabla_i T_{jp} - \nabla_j T_{ip}) \varphi_{lqk} g^{pq} g^{il} - T_{jp} g^{pq} T_{qk} + (\text{Tr } T) T_{jk} \\ & - T_{jl} T_{ab} g^{ap} g^{bq} \psi_{pqmk} g^{lm}. \end{aligned} \quad (4.55)$$

Proof. Equation (4.55) can be obtained from (4.53) by combining the first Bianchi identity of Riemannian geometry together with the identities of Theorem 4.23. The details can be found in [26, Section 4.2]. \square

Remark 4.56. Equation (4.55) shows that the metric of a torsion-free G_2 -structure is always *Ricci-flat*. (See also item (vi) of Remark 5.6 below.) \blacktriangle

On a general Riemannian manifold (M^n, g) , the Riemann curvature tensor R decomposes into the scalar curvature, the traceless Ricci curvature, and the conformally invariant *Weyl curvature*. When g is induced from a G_2 -structure φ , the Weyl tensor W decomposes further into three independent components W_{27} , W_{64} , and W_{77} as irreducible G_2 -representations. A detailed discussion of the curvature decomposition of G_2 -structures can be found in Cleyton–Ivanov [9] and in the forthcoming [13].

5 Exceptional Riemannian holonomy

In this section we briefly review the notion of the *holonomy* of a Riemannian manifold (M, g) , and place the geometry of torsion-free G_2 -structures in this context, as one of the geometries with *exceptional Riemannian holonomy*.

5.1 Parallel transport and Riemannian holonomy

Let (M^n, g) be a Riemannian manifold, and let ∇ be the Levi-Civita connection of the metric g . We review without proof the well-known basic properties of Riemannian holonomy. See, for example, [23, Chapters 2 & 3] for a more detailed discussion.

Definition 5.1. Fix $p \in M$. Let γ be loop based at p . This means that $\gamma : [0, 1] \rightarrow M$ is a continuous path, and piecewise smooth, such that $\gamma(0) = \gamma(1) = p$. Then, with respect to ∇ , the *parallel transport* $\Pi_\gamma : T_{\gamma(0)}M \rightarrow T_{\gamma(1)}M$ around the loop γ is a linear isomorphism of T_pM with itself, which depends on γ . We define the *holonomy* of the metric g at the point p , denoted $\text{Hol}_p(g)$, to be the set of all such isomorphisms. That is,

$$\text{Hol}_p(g) = \{\Pi_\gamma : T_pM \cong T_pM : \gamma \text{ is a loop based at } p\}.$$

It follows from the existence and uniqueness of parallel transport (which itself is a consequence of existence and uniqueness for systems of first order linear ordinary differential equations) that $\Pi_{\gamma\beta} = \Pi_\gamma \circ \Pi_\beta$, where

$\gamma \cdot \beta$ is the concatenation of paths, β followed by γ . Consequently, it is easy to see that $\text{Hol}_p(g)$ is closed under multiplication and inversion. That is, $\text{Hol}_p(g)$ is a *subgroup* of $\text{GL}(T_pM)$.

If we instead consider the restricted class of *contractible* loops at p , which is closed under concatenation of paths, we obtain the *restricted holonomy* of g at p , denoted $\text{Hol}_p^0(g)$. The group $\text{Hol}_p^0(g)$ is a normal subgroup of $\text{Hol}_p(g)$, and is the connected component of the identity. If M is simply-connected, then $\text{Hol}_p^0(g) = \text{Hol}_p(g)$ for all $p \in M$.

Because ∇ is the Levi-Civita connection, we have $\nabla g = 0$. Thus parallel transport with respect to ∇ preserves the inner product, and we conclude that in fact $\text{Hol}_p(g)$ is a subgroup of $\text{O}(T_pM, g_p)$, the group of isometries of the inner product space (T_pM, g_p) . Similarly $\text{Hol}_p^0(g)$ is a subgroup of $\text{SO}(T_pM, g_p)$, the group of orientation-preserving isometries of (T_pM, g_p) . \blacktriangle

The following proposition is straightforward to prove using the definitions.

Proposition 5.2. *The holonomy group $\text{Hol}_p(g)$ satisfies the following properties.*

- Let $p, q \in M$ lie in the same connected component of M . Then $\text{Hol}_p(g) \cong \text{Hol}_q(g)$. In fact, if γ is a piecewise smooth continuous path from p to q , and $P = \Pi_\gamma : T_pM \cong T_qM$ is the parallel transport isomorphism from T_pM to T_qM , then $\text{Hol}_q(g) = P \cdot \text{Hol}_p(g) \cdot P^{-1}$.
- Fix $p \in M$, and fix an isomorphism $T_pM \cong \mathbb{R}^n$. Then $\text{GL}(T_pM) \cong \text{GL}(n)$ and $\text{O}(T_pM, g_p) \cong \text{O}(n)$. With respect to this identification, $\text{Hol}_p(g)$ corresponds to a subgroup $H \subseteq \text{O}(n)$. If we choose any other isomorphism $T_pM \cong \mathbb{R}^n$, then the resulting subgroup \tilde{H} of $\text{O}(n)$ is in the same conjugacy class as H .
- Suppose M is connected. Then $\text{Hol}_p(g) \cong \text{Hol}_q(g)$ for all $p, q \in M$. Moreover, there exists a subgroup H of $\text{O}(n)$ such that $\text{Hol}_p(g) \cong H$ for all $p \in M$, and this subgroup H is unique up to conjugation.

Analogous statements hold for the restricted holonomy group $\text{Hol}_p^0(g)$, determining (when M is connected) a subgroup H^0 of $\text{SO}(n)$, unique up to conjugation.

Consequently, if M is connected, we abuse notation and call H the *holonomy group* and H^0 the *restricted holonomy group* of (M, g) . Observe that H and H^0 are not just abstract groups, but that they come naturally equipped with isomorphism classes of *representations on T_pM* for all $p \in M$.

Recall that a tensor S on M is called *parallel* if $\nabla S = 0$. There is a fundamental relationship between the holonomy group of g and the *parallel tensors* on M , given by the following.

Proposition 5.3. *Let (M, g) be a Riemannian manifold. Fix $p \in M$. Let $H \subseteq \text{GL}(T_pM)$ be the subgroup that leaves invariant $S|_p$ for all parallel tensors S on M .*

- We always have $\text{Hol}_p(g) \subseteq H$. Moreover, these two subgroups are usually equal. For example, this is the case if $\text{Hol}_p(g)$ is a closed subgroup of $\text{GL}(T_pM)$.
- If the group H fixes an element S_0 in some tensor space of T_pM , then there exists a parallel tensor S on M such that $S|_p = S_0$.

The way to think about Proposition 5.3 is as follows. The Riemannian holonomy H of a Riemannian manifold (M, g) is strictly smaller than $\text{O}(n)$ if and only if there exist nontrivial parallel tensors on M other than the metric g .

Remark 5.4. If M is simply-connected, then $H = H^0$ and consequently $H \subseteq \text{SO}(n)$. This means there exists a (necessarily parallel) Riemannian volume form $\mu \in \Omega^n(M)$ on M . This is consistent with the well-known fact from topology that any simply-connected manifold is orientable. \blacktriangle

5.2 The Berger classification of Riemannian holonomy

In 1955, Marcel Berger classified the possible Lie subgroups of $O(n)$ that could occur as Riemannian holonomy groups of a metric g , subject to the following technical hypotheses.

- We restrict to *simply-connected* manifolds. In general if (M, g) is not simply-connected then the holonomy H of (M, g) is a finite cover of the reduced holonomy H^0 . That is, the quotient H/H^0 is a discrete group.
- We must exclude the case when (M, g) is *locally reducible*. A locally reducible Riemannian manifold is *locally* a Riemannian product $(M_1, g_1) \times (M_2, g_2)$. In this case the Riemannian holonomy of (M, g) is a product of the holonomies of (M_1, g_1) and (M_2, g_2) .
- We must exclude the case when (M, g) is *locally symmetric*. A locally symmetric Riemannian manifold is *locally* isometric to a symmetric space $(G/H, g)$ where G is a group of isometries acting transitively on G/H with isotropy group H at any point. In this case the Riemannian holonomy of (M, g) is H .

Theorem 5.5 (Berger classification). *Let (M, g) be a simply-connected smooth Riemannian manifold of dimension n that is not locally reducible and not locally symmetric. Then the Riemannian holonomy $H \subseteq \text{SO}(n)$ can only be one of the following seven possibilities:*

$n = \dim M$	H	Parallel tensors	Name	Curvature
n	$\text{SO}(n)$	g, μ	<i>orientable</i>	
$2m$ ($m \geq 2$)	$\text{U}(m)$	g, ω	<i>Kähler</i>	
$2m$ ($m \geq 2$)	$\text{SU}(m)$	g, ω, Ω	<i>Calabi-Yau</i>	<i>Ricci-flat</i>
$4m$ ($m \geq 2$)	$\text{Sp}(m)$	$g, \omega_1, \omega_2, \omega_3, J_1, J_2, J_3$	<i>hyper-Kähler</i>	<i>Ricci-flat</i>
$4m$ ($m \geq 2$)	$(\text{Sp}(m) \times \text{Sp}(1))/\mathbb{Z}_2$	g, Υ	<i>quaternionic-Kähler</i>	<i>Einstein</i>
7	G_2	g, φ, ψ	G_2	<i>Ricci-flat</i>
8	$\text{Spin}(7)$	g, Φ	$\text{Spin}(7)$	<i>Ricci-flat</i>

Sketch of proof. Berger arrived at this classification by studying the *holonomy algebra* \mathfrak{h} of the holonomy group H . There is an intimate relation between \mathfrak{h} and the Riemann curvature operator $R \in \mathcal{S}^2(\mathfrak{so}(n))$ of g . First, because the Riemann curvature operator can be viewed as “infinitesimal holonomy”, it must be that $R \in \mathcal{S}^2(\mathfrak{h})$. Since it also satisfies the first Bianchi identity, this says that \mathfrak{h} cannot be too big. Second, by the Ambrose–Singer holonomy theorem, the span of the image of R at any point in M must generate \mathfrak{h} as a vector space, so \mathfrak{h} cannot be too small. Finally, for certain possible \mathfrak{h} , the fact that R must also satisfy the second Bianchi identity forces $\nabla R = 0$, in which case (M, g) is locally symmetric. Much more detailed discussion of this argument can be found in Joyce [23, Section 3.4]. \square

Remark 5.6. We make some remarks concerning the above table.

- The four restrictions $m \geq 2$ in the first column are mostly to eliminate redundancy, as we have the isomorphisms $\text{U}(1) \cong \text{SO}(2)$, $\text{Sp}(1) \cong \text{SU}(2)$, and $(\text{Sp}(1) \times \text{Sp}(1))/\mathbb{Z}_2 \cong \text{SO}(4)$. The case $\text{SU}(1)$ does not occur because $\text{SU}(1) \cong \{1\}$ is trivial and such a space is flat and thus symmetric.
- Because $\text{Sp}(k) \subseteq \text{SU}(2k) \subseteq \text{U}(2k)$, all hyper-Kähler manifolds are Calabi-Yau, and all Calabi-Yau manifolds are Kähler.
- Note that quaternionic-Kähler manifolds are in fact *not* Kähler. This ill-advised nomenclature has unfortunately stuck and is here to stay.
- Usually, the term *special holonomy* refers to any of the holonomy groups above other than the first two, perhaps because Kähler manifolds exist in sufficient abundance to not be that special.

- (v) The last two groups above, namely G_2 and $\text{Spin}(7)$, are called the *exceptional holonomy groups*. These Lie groups are both intimately related to the octonions \mathbb{O} . The connection between G_2 and \mathbb{O} is explained in Section 4.1 above. The connection between $\text{Spin}(7)$ and \mathbb{O} can be found, for example, in Harvey [17, Lemma 14.61] or Harvey–Lawson [18, Section IV.1.C.].
- (vi) The fact that metrics with special holonomy are all Einstein (including Ricci-flat) follows from consideration of the constraints on the Riemann curvature due to its relation with the holonomy algebra \mathfrak{h} , as explained in the sketch proof above. (See also Remark 4.56 above for the G_2 case.) \blacktriangle

It is interesting to note that Berger did not actually prove that all these groups *can actually occur* as Riemannian holonomy groups. He only excluded all other possibilities. It was widely suspected that the exceptional holonomies could not actually occur, only they could not be excluded using Berger’s method. We now know, of course, that *all* of the possibilities in the above table do occur, both in compact and in complete noncompact examples. See Section 6.2 for a brief survey of this history in the case of G_2 .

6 Torsion-free G_2 manifolds

In this section we discuss torsion-free G_2 manifolds, including a brief history of the search for irreducible examples, the known topological obstructions to existence in the compact case, and a comparison with Kähler and Calabi-Yau manifolds.

6.1 Irreducible and reducible torsion-free G_2 manifolds

Let (M, φ) be a torsion-free G_2 manifold. That is, φ is a torsion-free G_2 -structure as in Definition 4.43, and thus by Proposition 5.3 the holonomy $\text{Hol}(g_\varphi)$ of the induced Riemannian metric g_φ lies in G_2 .

Definition 6.1. We say (M, φ) is an *irreducible* torsion-free G_2 manifold if $\text{Hol}(g_\varphi) = G_2$. \blacktriangle

A torsion-free G_2 manifold could have *reduced holonomy*. That is, we could have $\text{Hol}(g_\varphi) \subsetneq G_2$. In fact there are some simple constructions that yield such reducible examples:

- If g_φ is flat, then $\text{Hol}(g_\varphi) = \{1\}$. In this case M is locally isomorphic to Euclidean \mathbb{R}^7 with the standard G_2 -structure φ_o .
- Let L^4 be a manifold with holonomy $\text{SU}(2) \cong \text{Sp}(1)$. This is a hyper-Kähler 4-manifold with hyper-Kähler triple $\omega_1, \omega_2, \omega_3$. Let X^3 be a flat Riemannian 3-manifold with global orthonormal parallel coframe e^1, e^2, e^3 . Let $M^7 = X^3 \times L^4$, and define a smooth 3-form φ on M by

$$\varphi = e^1 \wedge \omega_1 + e^2 \wedge \omega_2 + e^3 \wedge \omega_3 - e^1 \wedge e^2 \wedge e^3.$$

Then φ is a torsion-free G_2 -structure with $\text{Hol}(g_\varphi) = \text{SU}(2) \subsetneq G_2$. In this case we have

$$\psi = e^2 \wedge e^3 \wedge \omega_1 + e^3 \wedge e^1 \wedge \omega_2 + e^1 \wedge e^2 \wedge \omega_3 - \text{vol}_L$$

where $\text{vol}_L = \frac{1}{2}\omega_1^2 = \frac{1}{2}\omega_2^2 = \frac{1}{2}\omega_3^2$ is the volume form of L .

- Let L^6 be a manifold with holonomy $\text{SU}(3)$. This is a Calabi-Yau complex 3-fold with Kähler form ω and holomorphic volume form Ω . Let X^1 be a Riemannian 1-manifold with global unit parallel 1-form e^1 . Let $M^7 = X^1 \times L^6$, and define a smooth 3-form φ on M by

$$\varphi = e^1 \wedge \omega - \text{Re}\Omega.$$

Then φ is a torsion-free G_2 -structure with $\text{Hol}(g_\varphi) = \text{SU}(3) \subsetneq G_2$. In this case we have

$$\psi = \frac{1}{2}\omega^2 + e^1 \wedge \text{Im}\Omega.$$

Remark 6.2. If (M, φ) is a torsion-free G_2 manifold, then some criteria are known to determine if (M, φ) is irreducible. Here are two examples:

- (i) If M is *compact* with $\text{Hol}(g_\varphi) \subseteq G_2$, then $\text{Hol}(g_\varphi) = G_2$ if and only if the fundamental group $\pi_1(M)$ is *finite*. (See Joyce [23, Proposition 10.2.2].)
- (ii) If M is connected and *simply-connected*, with $\text{Hol}(g_\varphi) \subseteq G_2$, then $\text{Hol}(g_\varphi) = G_2$ if and only if there are *no nonzero parallel 1-forms*. (See Bryant–Salamon [5, Theorem 2].) ▲

6.2 A brief history of irreducible torsion-free G_2 manifolds

The search for examples of *irreducible* torsion-free G_2 manifolds (that is, Riemannian metrics with holonomy exactly G_2) has a long history. As explained in Section 5.2, it was originally believed such metrics could not exist. In this section we give a very brief and far from exhaustive survey of some of this history.

The first local (that is, incomplete) examples were found by Bryant [4] in 1987, using methods of exterior differential systems and Cartan–Kähler theory.

Then in 1989, Bryant–Salamon [5] found the first *complete noncompact* examples of G_2 holonomy metrics. These were metrics on the *total spaces of vector bundles*. Explicitly, these metrics were found on the bundles $\Lambda^2(S^4)$ and $\Lambda^2(\mathbb{C}P^2)$, which are rank 3 bundles over 4-dimensional bases, and on the bundle $\mathcal{S}(S^3)$, the spinor bundle of S^3 , which is a rank 4 bundle over a 3-dimensional base. These Riemannian manifold are all *asymptotically conical*. That is, the metrics approach Riemannian cone metrics at some particular rate at infinity. These torsion-free G_2 -structures are *cohomogeneity one*. That is, there is a Lie group of symmetries acting on (M, φ) with generic orbits of codimension one. Such symmetry reduces the partial differential equation $\nabla\varphi = 0$ to a (fully nonlinear) system of *ordinary differential equations*, which can be explicitly solved. The fact that the metrics have holonomy exactly G_2 was verified by using the criterion in item (ii) of Remark 6.2.

Remark 6.3. Since then, several explicit examples and a great many nonexplicit examples of complete noncompact holonomy G_2 metrics have been discovered, with various prescribed asymptotic geometry at infinity, such as asymptotically conical (AC), asymptotically locally conical (ALC), and others. In fact, very recent work of Foscolo–Haskins–Nordström [15, 16] has produced a spectacular new plethora of such examples. ▲

The first construction of *compact* irreducible torsion-free G_2 manifolds was given by Joyce [22] in 1994, and pushed further in the monograph [23]. The idea is the following. Start with the flat 7-torus T^7 , and take the quotient by a discrete group of isometries preserving the G_2 -structure φ_0 . The quotient is a singular *orbifold* with torsion-free G_2 -structure. Joyce then resolved the singularities by gluing in (quasi)-asymptotically locally Euclidean spaces with $SU(2)$ or $SU(3)$ holonomy, to produce a smooth compact 7-manifold M with *closed* G_2 -structure and “small” torsion. He then used analysis (see Section 7.1 below) to prove that M admits a torsion-free G_2 -structure. Finally, he showed the metrics had holonomy exactly G_2 by using the criterion (i) of Remark 6.2. This first construction is explained in more detail by Kovalev [30] in the present volume.

The second construction of *compact* irreducible torsion-free G_2 manifolds was introduced by Kovalev [29] in 2001 and pushed significantly further by Corti–Haskins–Nordström–Pacini [10] in 2015. It is called the “twisted connect sum construction”. The ideas is the following. Start with two noncompact *asymptotically cylindrical* Calabi–Yau complex 3-folds L_1 and L_2 , which are both asymptotic to $X^4 \times T^2$ where X^4 is a K3 complex surface. Take $L_1 \times S^1$ and $L_2 \times S^1$ and glue them together with a “twist” by identifying different factors of S^1 in order to obtain a smooth compact 7-manifold. The goal is then to construct a closed G_2 -structure on M with “small” torsion that can be perturbed using analysis to a torsion-free G_2 -structure (see Section 7.1 below). Being able to do this is a very delicate problem in algebraic geometry

involving “matching data”. This second construction is also explained in more detail by Kovalev [30] in the present volume.

More recently, a third construction of *compact* irreducible torsion-free G_2 manifolds appeared in Joyce–Karigiannis [24], involving glueing 3-dimensional families of Eguchi–Hanson spaces. This construction differs from the previous two because some of the noncompact “pieces” that are being glued together this time *do not* come equipped with torsion-free G_2 -structures. This is dealt with by solving a linear elliptic PDE on the noncompact Eguchi–Hanson space using weighted Sobolev spaces.

All three of the currently known constructions of compact irreducible torsion-free G_2 manifolds are similar in that they all use *glueing techniques* to construct a closed G_2 -structure φ with “small” torsion, and then invoke a general existence theorem of Joyce to prove that it can be *perturbed* to a nearby torsion-free G_2 -structure $\tilde{\varphi}$. This existence theorem is the subject of Section 7.1 below.

Thus, we know that Riemannian metrics with holonomy exactly G_2 *do exist* on compact manifolds, but they are not explicit. This is analogous to the case of Riemannian metrics with holonomy exactly $SU(m)$ (also called Calabi–Yau metrics) on compact manifolds. By Yau’s proof of the Calabi conjecture, we know that many such metrics exist, but we cannot describe them explicitly. In fact, special holonomy metrics on compact manifolds should in some sense be thought of as “transcendental” objects.

So far we have only found G_2 -holonomy metrics that are “close to the edge of the moduli space”. That is, these metrics are close to either developing singularities or tearing apart into two disjoint noncompact pieces. That is, the three known constructions of compact irreducible torsion-free G_2 manifolds are very likely producing only a very small part of the “landscape” of holonomy G_2 metrics.

6.3 Cohomological obstructions to existence in the compact case

There are several known *cohomological obstructions* to the existence of torsion-free G_2 -structures on a compact manifold. We describe some of these in this section. Let (M, φ) be a *compact* manifold with a *torsion-free* G_2 -structure φ . Let g_φ be the Riemannian metric induced by φ . Thus $\text{Hol}(g_\varphi) \subseteq G_2$. Since (M, g_φ) is a compact oriented Riemannian manifold, the Hodge Theorem applies. That is, any deRham cohomology class has a unique harmonic representative.

Since φ is torsion-free, by Corollary 4.50, the form φ is closed and coclosed and thus harmonic. Because $\varphi \neq 0$, we deduce from the Hodge Theorem that $[\varphi]$ is a nontrivial class in $H^3(M, \mathbb{R})$. Hence we find our first cohomological obstruction:

$$b^3 \geq 1 \quad \text{if } M \text{ admits a torsion-free } G_2\text{-structure.}$$

where $b^k = \dim H^k(M, \mathbb{R})$ is the k^{th} Betti number of M . The same argument applies to ψ , so $b^4 \geq 1$, but $b^4 = b^3$ by Poincaré duality, so this is not new information.

Suppose $\text{Hol}(g_\varphi) = G_2$. Then by item (i) of Remark 6.2 we must have $\pi_1(M)$ is finite. It follows from algebraic topology that $H^1(M, \mathbb{R}) = \{0\}$. Hence we find our second cohomological obstruction:

$$b^1 = 0 \quad \text{if } M \text{ admits an } \textit{irreducible} \text{ torsion-free } G_2\text{-structure.} \tag{6.4}$$

Before we can discuss the two other cohomological obstructions, we need to explain the interaction of the representation-theoretic decompositions of Section 4.3 with the Hodge Theorem.

Because φ is torsion-free, one can show that the Hodge Laplacian Δ_d *commutes* with the orthogonal projection operators onto the irreducible summands of the decomposition of Ω^\bullet described in Section 4.3. (See Joyce [23, Theorem 3.5.3] for details.) Combining this fact with the Hodge Theorem, we conclude that the decompositions of Section 4.3 *descend to deRham cohomology*. That is, if we define

$$\mathcal{H}_l^k = \{\gamma \in \Omega_l^k \mid \Delta_d \gamma = 0\}$$

to be the space of *harmonic* Ω_l^k -forms, and \mathcal{H}^k to be the space of harmonic k -forms, then we have

$$\begin{aligned}\mathcal{H}^2 &= \mathcal{H}_7^2 \oplus \mathcal{H}_{14}^2, \\ \mathcal{H}^3 &= \mathcal{H}_1^3 \oplus \mathcal{H}_7^3 \oplus \mathcal{H}_{27}^3.\end{aligned}\tag{6.5}$$

Moreover, it follows from the explicit descriptions of Ω_l^k in Section 4.3 and the fact that Δ_d commutes with the projections and with the Hodge star \star that

$$\Delta_d(f\varphi) = (\Delta_d f)\varphi, \quad \Delta_d(\alpha \wedge \varphi) = (\Delta_d \alpha) \wedge \varphi,$$

for all $f \in \Omega^0$ and all $\alpha \in \Omega^1$. These identities imply that

$$\begin{aligned}\mathcal{H}_1^0 &\cong \mathcal{H}_1^3 \cong \mathcal{H}_1^4 \cong \mathcal{H}_1^7, & \mathcal{H}_7^1 &\cong \mathcal{H}_7^2 \cong \mathcal{H}_7^3 \cong \mathcal{H}_7^4 \cong \mathcal{H}_7^5 \cong \mathcal{H}_7^6, \\ \mathcal{H}_{14}^2 &\cong \mathcal{H}_{14}^5, & \mathcal{H}_{27}^3 &\cong \mathcal{H}_{27}^4.\end{aligned}$$

Define $b_l^k = \dim \mathcal{H}_l^k$ to be the ‘‘refined Betti numbers’’ of (M, φ) . Then we have shown that

$$b_l^k = b_{l'}^{k'} \quad \text{if } l = l'.$$

In particular $b_7^k = b_7^1 = b^1$ for $k = 2, \dots, 6$, and $b_1^k = b_1^0 = b^0$ for $k = 3, 4, 7$. We deduce that

$$\begin{aligned}b^2 &= b_7^2 + b_{14}^2 = b^1 + b_{14}^2, \\ b^3 &= b_1^3 + b_7^3 + b_{27}^3 = b^0 + b^1 + b_{27}^3.\end{aligned}$$

(Note that if M is connected then $b^0 = 1$, and if in addition φ is irreducible then by (6.4) we get $b^2 = b_{14}^2$ and $b^3 = 1 + b_{27}^3$.)

There exists a real quadratic form Q on $H^2(M, \mathbb{R})$ defined as follows. Let $[\beta] \in H^2(M, \mathbb{R})$. Define

$$Q([\beta]) = \int_M \beta \wedge \beta \wedge \varphi.\tag{6.6}$$

(In fact, it is easy to see using Stokes’s Theorem that Q is well-defined as long as $d\varphi = 0$, and that Q depends only on the *cohomology class* $[\varphi] \in H^3(M, \mathbb{R})$. We do not need torsion-free to define Q .)

But now suppose that φ is not only torsion-free, but also irreducible. Then by (6.4) and the discussion above, we have $\mathcal{H}^2 = \mathcal{H}_{14}^2$. Given a cohomology class $[\beta] \in H^2(M, \mathbb{R})$, the Hodge theorem gives us a unique harmonic representative β_H , which must lie in Ω_{14}^2 . By (4.31), we have $\beta_H \wedge \varphi = \star \beta_H$, and hence

$$Q([\beta]) = \int_M \beta_H \wedge \beta_H \wedge \varphi = \int_M \beta_H \wedge \star \beta_H = \int_M \|\beta_H\|^2 \text{vol} \geq 0$$

with equality if and only if $\beta_H = 0$, which is equivalent to $[\beta] = 0$. Hence we find our third cohomological obstruction:

- Let φ be a *closed* G_2 -structure on a compact manifold M with $\pi_1(M)$ finite. (So that any torsion-free G_2 -structure on M must necessarily be irreducible.) If there exists a torsion-free G_2 -structure in the cohomology class $[\varphi] \in H^3(M, \mathbb{R})$, then the quadratic form Q defined in (6.6) must be *positive definite*.

Remark 6.7. Recall Remark 4.37. If the other orientation is used, then Q must be *negative definite*. So we can unambiguously state the third cohomological obstruction as saying that Q must be *definite*. Moreover, if φ is merely torsion-free but not irreducible, it is easy to see from (4.31) that (with our convention for the orientation), the *signature* of Q is $(b^2 - b^1, b^1)$. \blacktriangle

Finally, recall from Chern–Weil theory that a compact 7-manifold M has a *real first Pontryagin class* $p_1(TM) \in H^4(M, \mathbb{R})$, defined as the cohomology class represented by the closed 4-form $\frac{1}{8\pi^2} \text{Tr}(R \wedge R)$

where $R \in \Gamma(\text{End}(TM) \otimes \Lambda^2 T^*M)$ is the curvature form of *any* connection on TM . If φ is torsion-free, then g_φ has holonomy contained in G_2 , and hence, because Riemann curvature is “infinitesimal holonomy” we have that in fact $R \in \Gamma(\text{End}(TM) \otimes \Lambda_{14}^2 T^*M)$. That is, the 2-form part of R lies in Ω_{14}^2 . But then by (4.31) we have

$$\text{Tr}(R \wedge R) \wedge \varphi = \text{Tr}(R \wedge *R) = |R|^2 \text{vol},$$

and thus

$$(p_1(TM) \cup [\varphi]) \cdot [M] = \frac{1}{8\pi^2} \int_M \text{Tr}(R \wedge R) \wedge \varphi = \frac{1}{8\pi^2} \int_M |R|^2 \text{vol},$$

where $[M] \in H_7(M)$ is the fundamental class of M and \cdot denotes the canonical pairing between $H^7(M, \mathbb{R})$ and $H_7(M)$. This is clearly positive unless R is identically zero. Hence we find our fourth cohomological obstruction:

$$p_1(TM) \neq 0 \quad \text{if } M \text{ admits a } \textit{nonflat} \text{ torsion-free } G_2\text{-structure.}$$

6.4 Comparison with Kähler and Calabi-Yau manifolds

In this section we make some remarks about the similarities and the differences between torsion-free G_2 manifolds and Kähler manifolds in general and Calabi-Yau manifolds in particular. A good reference for Kähler and Calabi-Yau geometry is Huybrechts [21].

Manifolds with $U(m)$ -structure are in some ways analogous to manifolds with G_2 -structure, as detailed in the following table.

	$U(m)$ -structure on (M^{2m}, g)	G_2 -structure on (M^7, g)
Nondegenerate form	$\omega \in \Omega^2$	$\varphi \in \Omega^3$
Vector cross product	$J \in \Gamma(TM \otimes T^*M)$	$\times \in \Gamma(TM \otimes \Lambda^2 T^*M)$
Fundamental relation	$\omega(u, v) = g(Ju, v)$	$\varphi(u, v, w) = g(u \times v, w)$

One very important difference between $U(m)$ -structures and G_2 -structures was already mentioned in Remark 4.10, but it is so crucial that it is worth repeating here. For a $U(m)$ -structure, the metric g and the nondegenerate 2-form ω are essentially independent, subject only to mild compatibility conditions, and together they determine J . In contrast, for a G_2 -structure the nondegenerate 3-form φ *determines* the metric g and consequently the cross product \times as well.

Now consider the *torsion-free* cases of such structures. A $U(m)$ -structure is torsion-free if $\nabla\omega = 0$. Such manifolds are called *Kähler* and have Riemannian holonomy contained in the Lie subgroup $U(m)$ of $SO(2m)$. A G_2 -structure is torsion-free if $\nabla\varphi = 0$. Such manifolds have Riemannian holonomy contained in the Lie subgroup G_2 of $SO(7)$. In the torsion-free cases, both ω and φ are *calibrations*. (See [35, 34] in the present volume for more about calibrations.) Both Kähler manifolds and torsion-free G_2 manifolds also admit special *connections* on vector bundles, namely the Hermitian-Yang-Mills connections and the G_2 -instantons, respectively. (See [41] in the present volume for more about G_2 -instantons.)

Here is where we see another very important difference. As we saw in Remark 4.56, the metric g_φ of a torsion-free G_2 -structure is always Ricci-flat. But the metric g of a Kähler manifold need *not* be Ricci-flat. In fact, the Calabi–Yau Theorem, gives a topological characterization (in the compact case) of exactly which Kähler metrics are Ricci-flat. They are precisely those metrics with holonomy contained in the Lie subgroup $SU(m)$ of $U(m)$. The precise statement of the Calabi–Yau theorem is as follows.

Theorem 6.8. *Let M be a compact Kähler manifold, with Kähler form ω . Then there exists a Ricci-flat Kähler metric $\tilde{\omega}$ in the same cohomology class as ω if and only if $c_1(TM) = 0$, where $c_1(TM)$ is the first Chern class of TM . Moreover, when it exists the Ricci-flat Kähler metric is unique.*

We are very far from having an analogous theorem in G_2 geometry. In fact, we do not even have any idea of what the correct conjecture might be. The main tool that allowed Yau to reformulate the Calabi conjecture into a statement about existence and uniqueness of solutions to a complex Monge-Ampère

equation is the $\partial\bar{\partial}$ -lemma in Kähler geometry. There is no close analogue of this result for torsion-free G_2 manifolds.

Heuristically, the Calabi–Yau Theorem allows us to go from $U(m)$ holonomy to $SU(m)$ holonomy, which is a reduction in the dimension of the holonomy group from m^2 to $m^2 - 1$, a difference of 1 dimension, and it corresponds to an (albeit fully nonlinear) *scalar* partial differential equation. In contrast, to obtain a Riemannian metric with holonomy G_2 , we must start with $SO(7)$ holonomy. Thus we need to reduce the dimension of the holonomy group from 21 to 14, so we expect a system of 7 *equations*, or equivalently a single partial differential equation for an unknown *1-form* rather than for an unknown function as in the Calabi–Yau Theorem. Precisely how such a heuristic discussion can be made into a precise mathematical conjecture remains a mystery at present.

In fact, a better analogy is the following. Let M^{2m} be a compact manifold that admits $U(m)$ -structures. What are *necessary and sufficient* topological conditions that ensure that M^{2m} admits a Kähler structure? We know many necessary conditions. (See Huybrechts [21], for example.) But we are very far from knowing sufficient conditions.

7 Three theorems about compact torsion-free G_2 -manifolds

In this final section we briefly discuss three important theorems about compact torsion-free G_2 manifolds: an existence theorem of Joyce, the smoothness of the moduli space (also due to Joyce), and a variational characterization of compact torsion-free G_2 manifolds due to Hitchin. Only the main ideas of the proofs are sketched. We refer the reader to the original sources for the details.

7.1 An existence theorem for compact torsion-free G_2 manifolds

In Section 6.2 we discussed known constructions of compact irreducible torsion-free G_2 manifolds. These constructions invoke the only analytic existence theorem that is known for torsion-free G_2 -structures, which is a result of Joyce that originally appeared in [22] but which can also be found in [23, Section 11.6]. As mentioned in Section 6.2, the idea is that if one has a *closed* G_2 -structure φ on M whose torsion is sufficiently small, the theorem guarantees the existence of a “nearby” torsion-free G_2 -structure $\tilde{\varphi}$ that is in the same cohomology class as φ . The statement of the theorem that we give here is a slightly modified version given in [24, Theorem 2.7].

Theorem 7.1 (Existence Theorem of Joyce). *Let α , K_1 , K_2 , and K_3 be any positive constants. Then there exist $\varepsilon \in (0, 1]$ and $K_4 > 0$, such that whenever $0 < t \leq \varepsilon$, the following holds.*

Let (M, φ) be a compact 7-manifold with G_2 -structure φ satisfying $d\varphi = 0$. Suppose there exists a closed 4-form η such that

- (i) $\|\star_\varphi\varphi - \eta\|_{C^0} \leq K_1 t^\alpha$,
- (ii) $\|\star_\varphi\varphi - \eta\|_{L^2} \leq K_1 t^{\frac{7}{2} + \alpha}$,
- (iii) $\|d(\star_\varphi\varphi - \eta)\|_{L^{14}} \leq K_1 t^{-\frac{1}{2} + \alpha}$,
- (iv) the injectivity radius inj of g_φ satisfies $\text{inj} \geq K_2 t$,
- (v) the Riemann curvature tensor Rm of g_{ph} satisfies $\|\text{Rm}\|_{C^0} \leq K_3 t^{-2}$.

Then there exists a smooth torsion-free G_2 -structure $\tilde{\varphi}$ on M such that $\|\tilde{\varphi} - \varphi\|_{C^0} \leq K_4 t^\alpha$ and $[\tilde{\varphi}] = [\varphi]$ in $H^3(M, \mathbb{R})$. Here all norms are computed using the original metric g_φ .

We make some remarks about the conditions (i)–(iii) of the theorem. Since φ is closed, it would be torsion-free if and only if $\star_\varphi\varphi$ were also closed. The hypotheses (i)–(iii) above say that $\star_\varphi\varphi$ is *almost*

closed, in that there exists a closed 4-form η that is close to $\star_\varphi\varphi$ in various norms, namely the C^0 , L^2 , and (essentially) the $W^{14,1}$ norms.

The idea of the proof of Theorem 7.1 is as follows. Since we want $\tilde{\varphi}$ is to be in the same cohomology class as φ , we must have $\tilde{\varphi} = \varphi + d\sigma$ for some $\sigma \in \Omega^2$, and by Hodge theory we can assume that $d^*\sigma = 0$. Joyce shows that the torsion-free condition

$$d(\star_{\varphi+d\sigma}(\varphi + d\sigma)) = 0$$

can be rewritten as

$$\Delta_d\sigma = \mathcal{Q}(\sigma, d\sigma) \tag{7.2}$$

where \mathcal{Q} is some nonlinear expression that is at least order two in $d\sigma$. Joyce shows that the above equation can be solved by iteration. Explicitly, taking $\sigma_0 = 0$, then for each $k \geq 1$, Joyce solves the series of linear equations

$$\Delta_d\sigma_k = \mathcal{Q}(\sigma_{k-1}, d\sigma_{k-1}).$$

Using the *a priori estimates (i)–(iii)*, Joyce then shows that $\lim_{k \rightarrow \infty} \sigma_k$ exists as a smooth 2-form satisfying (7.2). This is essentially a “fixed-point theorem” type of argument. The complete details can be found in [23, Section 11.6].

7.2 The moduli space of compact torsion-free G_2 -structures

Whenever one studies a certain type of structure in mathematics, it is natural to consider the “set of all possible such structures”, modulo a reasonable notion of equivalence. Usually this “moduli space” of structures has its own special structure, and an understanding of the special structure on the moduli space sometimes yields information about the original object on which such structures are defined.

In our setting, consider a compact torsion-free G_2 manifold (M, φ) . We want to consider the set of all possible torsion-free G_2 -structures on the same underlying smooth 7-manifold M , modulo a reasonable notion of equivalence. The usual notion of equivalence in differential geometry is *diffeomorphism*. Indeed, if φ is a torsion-free G_2 -structure on M and $F : M \rightarrow M$ is a diffeomorphism, then it is easy to see that $F^*\varphi$ is also a torsion-free G_2 -structure on M , with metric $g_{F^*\varphi} = F^*g_\varphi$.

In fact, it is more convenient to consider only those diffeomorphisms of M that are *isotopic to the identity*. That is, those diffeomorphisms that are connected to the identity map on M by a continuous path in the space Diff of diffeomorphisms of M . This is the *connected component of the identity* in Diff , and we denote it by Diff_0 . The reason we prefer the space Diff_0 is because *it acts trivially on cohomology*. That is, suppose $[\alpha] \in H^k(M, \mathbb{R})$ and let $F \in \text{Diff}_0$. Then we claim that $[F^*\alpha] = [\alpha]$. To see this, let F_t be a continuous path in Diff with $F_0 = \text{Id}_M$ and $F_1 = F$, given by the flow of the vector field X_t on M . Since α is a closed form, we have

$$\begin{aligned} F^*\alpha - \alpha &= \int_0^1 \frac{d}{dt}(F_t^*\alpha) = \int_0^1 \mathcal{L}_{X_t}\alpha \\ &= \int_0^1 (dX_t \lrcorner \alpha + X_t \lrcorner d\alpha) = \int_0^1 dX_t \lrcorner \alpha \\ &= d\left(\int_0^1 X_t \lrcorner \alpha\right), \end{aligned}$$

and thus $F^*\alpha - \alpha$ is exact.

Definition 7.3. Let (M, φ_0) be a compact torsion-free G_2 manifold. Let \mathcal{T} be the set of all torsion-free G_2 -structures on M . That is,

$$\mathcal{T} = \{\varphi \in \Omega_+^3 \mid d\varphi = 0, d\star_\varphi\varphi = 0\}.$$

The *moduli space* \mathcal{M} of torsion-free G_2 -structures on M is defined to be the quotient topological space

$$\mathcal{M} = \mathcal{T}/\text{Diff}_0$$

of \mathcal{T} by the action of Diff_0 . ▲

Remark 7.4. The space \mathcal{M} in Definition 7.3 should probably more properly be called the *Teichmüller space*, and then the “moduli space” would be the quotient \mathcal{T}/Diff by the full diffeomorphism group, in analogy with the usage of terminology for Riemann surfaces. However, the nomenclature we have given in Definition 7.3 is standard in the field of G_2 geometry. ▲

The first important result that was established about the moduli space was the following theorem of Joyce, that originally appeared in [22] but which can also be found in [23, Section 10.4].

Theorem 7.5 (Moduli Space Theorem of Joyce). *Let M be a compact 7-manifold with torsion-free G_2 -structure φ_0 . The moduli space \mathcal{M} of torsion-free G_2 -structures on M is a smooth manifold of dimension $b^3 = \dim H^3(M, \mathbb{R})$. In fact, the “period map” $\mathcal{P} : \mathcal{M} \rightarrow H^3(M, \mathbb{R})$ that takes an equivalence class $[\varphi]_{\mathcal{M}}$ in the quotient space $\mathcal{M} = \mathcal{T}/\text{Diff}_0$ to the deRham cohomology class $[\varphi]$ is a local diffeomorphism.*

The idea of the proof of Theorem 7.5 is as follows. Joyce constructs a “slice” \mathcal{S}_φ for the action of Diff_0 on \mathcal{T} in a neighbourhood of any $\varphi \in \mathcal{T}$. A slice \mathcal{S}_φ is a submanifold of \mathcal{T} containing φ that is locally transverse to the orbits of Diff_0 near φ . This means that all nearby orbits of Diff_0 each intersect \mathcal{T} at only one point. Then $\mathcal{M} = \mathcal{T}/\text{Diff}_0$ is locally homeomorphic in a neighbourhood of $[\varphi]_{\mathcal{M}} \in \mathcal{M}$ to \mathcal{S}_φ . Since $\varphi \in \mathcal{T}$ is arbitrary, we deduce that \mathcal{M} is a smooth manifold of dimension $\dim \mathcal{S}$.

In fact a slice \mathcal{S}_φ is given by

$$\mathcal{S}_\varphi = \{\tilde{\varphi} \in \Omega_+^3 \mid d\tilde{\varphi} = 0, d \star_{\tilde{\varphi}} \tilde{\varphi} = 0, \pi_7(d^* \tilde{\varphi}) = 0\}, \quad (7.6)$$

where π_7 is the orthogonal projection $\pi_7 : \Omega^2 \rightarrow \Omega_7^2$ with respect to the G_2 -structure φ . The way to understand where the above \mathcal{S}_φ comes from is to consider tangent vectors to the orbit of Diff_0 at φ . Such a tangent vector is of the form

$$\left. \frac{d}{dt} \right|_{t=0} h_t^* \varphi = \mathcal{L}_X \varphi = d(X \lrcorner \varphi)$$

where h_t is the flow of a smooth vector field X on M . By the description (4.31), the tangent space at φ of the orbit of Diff_0 is thus the space $d(\Omega_7^2)$. It thus makes sense to define

$$\mathcal{S}_\varphi = \{\tilde{\varphi} \in \mathcal{T} \mid \langle \tilde{\varphi} - \varphi, d(X \lrcorner \varphi) \rangle_{L^2} = 0 \forall X \in \Gamma(TM)\}, \quad (7.7)$$

because for $\tilde{\varphi}$ close to φ , the condition of L^2 -orthogonality to the tangent spaces of the orbit of Diff_0 through φ would ensure local transversality. Since φ is torsion-free, we have $d^* \varphi = 0$. Thus integration by parts shows that (7.7) is equivalent to (7.6).

It still remains to explain why \mathcal{S}_φ is a smooth manifold of dimension b^3 . Given $\tilde{\varphi} \in \mathcal{T}$, by Hodge theory with respect to g_φ we can write $\tilde{\varphi} = \varphi + \xi + d\eta$ for some $\xi \in \mathcal{H}^3$ and some $\eta \in d^*(\Omega^3)$. For $\tilde{\varphi}$ sufficiently close to φ in the C^0 norm, Joyce shows that

$$\tilde{\varphi} \in \mathcal{S}_\varphi \iff \Delta_d \eta = \star d(\mathcal{Q}(\xi, d\eta)) \quad (7.8)$$

where \mathcal{Q} is a nonlinear expression that is at least order two in $d\eta$. This is a fully nonlinear elliptic equation for η given any $\xi \in \mathcal{H}^3$. Using the Banach Space Implicit Function Theorem, Joyce shows that the space of solutions (ξ, η) to the right hand side of (7.8) is a smooth manifold of dimension b^3 . The complete details can be found in [23, Section 10.4].

Remark 7.9. A consequence of the fact from Theorem 7.5 that the period map $\mathcal{P} : \mathcal{M} \rightarrow H^3(M, \mathbb{R})$ is a local diffeomorphism is the following. The manifold \mathcal{M} has a natural *affine structure*, that is a covering by coordinate charts whose transition functions are affine maps. In Karigiannis–Leung [28] this affine structure is exploited to study special structures on \mathcal{M} , including a natural *Hessian metric* and a *symmetric cubic form* called the “Yukawa coupling”. This Hessian metric is obtained from the *Hitchin volume functional* defined in Section 7.3 below. ▲

We know *very little* about the *global structure* of \mathcal{M} . (But see the survey article by Crowley–Goette–Nordström [11] in the present volume for some recent progress on the (dis-)connectedness of \mathcal{M} .)

7.3 A variational characterization of torsion-free G_2 -structures

It is the case that some natural geometric structures can be given a *variational interpretation*. That is, they can be characterized as critical points of a certain natural geometric functional, which means that they are solutions to the associated Euler-Lagrange equations for this functional. Some examples of such geometric structures and their associated functionals are:

- minimal submanifolds (the volume functional),
- harmonic maps (the energy functional),
- Einstein metrics (the Einstein-Hilbert functional),
- Yang-Mills connections (the Yang-Mills functional).

It was an important observation of Hitchin [19] that torsion-free G_2 -structures on compact manifolds can be given a variational interpretation. The setup is as follows. Let M^7 be compact, and as usual, let Ω_+^3 be the set of G_2 -structures on M . Given $\varphi \in \Omega_+^3$, we get a metric g_φ , a Riemannian volume vol_φ , and a dual 4-form $\star_\varphi \varphi$.

Definition 7.10. The *Hitchin functional* is defined to be the map $\mathcal{F} : \Omega_+^3 \rightarrow \mathbb{R}$ given by

$$\mathcal{F}(\varphi) = \int_M \varphi \wedge \star_\varphi \varphi = 7 \int_M \text{vol}_\varphi = 7 \text{Vol}(M, g_\varphi), \quad (7.11)$$

where we have used the fact that $|\varphi|_{g_\varphi}^2 = 7$ from (4.2). Thus, up to a positive factor, $\mathcal{F}(\varphi)$ is the total volume of M with respect to the metric g_φ . \blacktriangle

Hitchin's observation was to restrict \mathcal{F} to a *cohomology class* containing a closed G_2 -structure. That is, suppose φ_0 is a *closed* G_2 -structure on M , and let

$$\mathcal{C}_\varphi = \Omega_+^3 \cap [\varphi] = \{\tilde{\varphi} \in \Omega_+^3 \mid d\tilde{\varphi} = 0, [\tilde{\varphi}] = [\varphi] \in H^3(M, \mathbb{R})\}.$$

In [19], Hitchin proved the following.

Theorem 7.12 (Hitchin's variational characterization). *Let φ be a closed G_2 -structure on M , and consider the restriction of \mathcal{F} to the set \mathcal{C}_φ defined above. Then φ is a critical point of $\mathcal{F}|_{\mathcal{C}_\varphi}$ if and only if φ is torsion-free. That is,*

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{F}(\varphi + t d\eta) = 0 \iff d \star_\varphi \varphi = 0.$$

Moreover, at a critical point φ , the second variation of $\mathcal{F}|_{\mathcal{C}_\varphi}$ is nonpositive. This means that critical points are local maxima.

The proof of Theorem 7.12 is quite straightforward given the following observation, which is quite useful itself in many other applications. Let $\varphi(t)$ be a smooth family of G_2 -structures with $\left. \frac{d}{dt} \right|_{t=0} \varphi(t) = \gamma$. Then

$$\left. \frac{d}{dt} \right|_{t=0} \star_{\varphi(t)} \varphi(t) = \frac{4}{3} \star \pi_1 \gamma + \star \pi_7 \gamma - \star \pi_{27} \gamma, \quad (7.13)$$

where the orthogonal projections $\pi_k : \Omega^3 \rightarrow \Omega_k^3$ and the Hodge star \star are all taken with respect to $\varphi(0)$. Two different proofs of (7.13) can be found in [19] and in [26, Remark 3.6].

The interesting observation in Theorem 7.12 that torsion-free G_2 -structures are *local maxima* of \mathcal{F} restricted to a cohomology class motivates the idea to try to *flow* to a torsion-free G_2 -structure by taking the appropriate *gradient flow* of \mathcal{F} . This yields the *Laplacian flow* of closed G_2 -structures. See the article by Lotay [36] in the present volume for a discussion of geometric flows of G_2 -structures, including the Laplacian flow.

References

- [1] I. Agricola, “Old and new on the exceptional group G_2 ”, *Notices Amer. Math. Soc.* **55** (2008), 922–929. MR2441524
- [2] I. Agricola, S. Chiossi, T. Friedrich, and J. Höll, “Spinorial description of $SU(3)$ - and G_2 -manifolds”, *J. Geom. Phys.* **98** (2015), 535–555. MR3414976
- [3] R. B. Brown and A. Gray, “Vector cross products”, *Comment. Math. Helv.* **42** (1967), 222–236. MR0222105
- [4] R.L. Bryant, “Metrics with exceptional holonomy”, *Ann. of Math. (2)* **126** (1987), 525–576. MR0916718
- [5] R.L. Bryant and S.M. Salamon, “On the construction of some complete metrics with exceptional holonomy”, *Duke Math. J.* **58** (1989), 829–850. MR1016448
- [6] R.L. Bryant, “Some remarks on G_2 -structures”, *Proceedings of Gökova Geometry-Topology Conference 2005*, 75–109. MR2282011 <https://arxiv.org/abs/math/0305124>
- [7] D.R. Cheng, S. Karigiannis, and J. Madnick, “Bubble Tree Convergence of Conformally Cross Product Preserving Maps”, *submitted for publication*. <https://arxiv.org/abs/1909.03512>
- [8] S. Chiossi and S. Salamon, “The intrinsic torsion of $SU(3)$ and G_2 structures”, in *Differential geometry, Valencia, 2001*, 115–133, World Sci. Publ., River Edge, NJ. MR1922042
- [9] R. Cleyton and S. Ivanov, “Curvature decomposition of G_2 -manifolds”, *J. Geom. Phys.* **58** (2008), 1429–1449. MR2453675
- [10] A. Corti, M. Haskins, J. Nordström, and T. Pacini, “ G_2 -manifolds and associative submanifolds via semi-Fano 3-folds”, *Duke Math. J.* **164** (2015), 1971–2092. MR3369307
- [11] D. Crowley, S. Goette, and J. Nordström, “Distinguishing G_2 -manifolds”, *Lectures and Surveys on G_2 -manifolds and related topics*, Fields Institute Communications, Springer. (The present volume.)
- [12] X. de la Ossa, S. Karigiannis, and E. Svanes, “Geometry of $U(m)$ -structures: Kähler identities, the dd^c lemma, and Hodge theory”, *in preparation*.
- [13] S. Dwivedi, P. Gianniotis, and S. Karigiannis, “Flows of G_2 -structures, II: Curvature, torsion, symbols, and functionals”; *in preparation*.
- [14] M. Fernández and A. Gray, “Riemannian manifolds with structure group G_2 ”, *Ann. Mat. Pura Appl. (4)* **132** (1982), 19–45. MR0696037
- [15] L. Foscolo, M. Haskins, and J. Nordström, “Complete non-compact G_2 -manifolds from asymptotically conical Calabi-Yau 3-folds”, <https://arxiv.org/abs/1709.04904>
- [16] L. Foscolo, M. Haskins, and J. Nordström, “Infinitely many new families of complete cohomogeneity one G_2 -manifolds: G_2 analogues of the Taub-NUT and Eguchi-Hanson spaces”, <https://arxiv.org/abs/1805.02612>
- [17] R. Harvey, *Spinors and calibrations*, Perspectives in Mathematics, 9, Academic Press, Inc., Boston, MA, 1990. MR1045637
- [18] R. Harvey and H.B. Lawson, “Calibrated geometries”, *Acta Math.* **148** (1982), 47–157. MR0666108
- [19] N. Hitchin, “The geometry of three-forms in six and seven dimensions”; <https://arxiv.org/abs/math/0010054>
- [20] N. Hitchin, “Stable forms and special metrics”, in *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, 70–89, *Contemp. Math.* **288**, Amer. Math. Soc., Providence, RI. MR1871001
- [21] D. Huybrechts, *Complex geometry*, Universitext, Springer-Verlag, Berlin, 2005. MR2093043
- [22] D.D. Joyce, “Compact Riemannian 7-manifolds with holonomy G_2 . I, II”, *J. Differential Geom.* **43** (1996), 291–328, 329–375. MR1424428

- [23] D.D. Joyce, *Compact manifolds with special holonomy*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000. MR1787733
- [24] D. Joyce and S. Karigiannis, “A new construction of compact torsion-free G_2 manifolds by gluing families of Eguchi-Hanson spaces”, *J. Differential Geom.*, to appear. <https://arxiv.org/abs/1707.09325>
- [25] S. Karigiannis, “Deformations of G_2 and $\text{Spin}(7)$ structures on manifolds”, *Canad. J. Math.* **57** (2005), 1012–1055. MR2164593
- [26] S. Karigiannis, “Flows of G_2 -structures, I.”, *Q. J. Math.* **60** (2009), 487–522. MR2559631 <http://arxiv.org/abs/math/0702077>
- [27] S. Karigiannis, “Some notes on G_2 and $\text{Spin}(7)$ geometry”, *Recent Advances in Geometric Analysis; Advanced Lectures in Mathematics*, Vol. 11; International Press, (2010), 129–146. <https://arxiv.org/abs/math/0608618>
- [28] S. Karigiannis and N.C. Leung, “Hodge theory for G_2 -manifolds: intermediate Jacobians and Abel-Jacobi maps”, *Proc. Lond. Math. Soc. (3)* **99** (2009), 297–325. MR2533667
- [29] A. Kovalev, “Twisted connected sums and special Riemannian holonomy”, *J. Reine Angew. Math.* **565** (2003), 125–160. MR2024648
- [30] A. Kovalev, “Constructions of compact G_2 -holonomy manifolds”, *Lectures and Surveys on G_2 -manifolds and related topics*, Fields Institute Communications, Springer. (The present volume.)
- [31] H.B. Lawson, Jr. and M.-L. Michelsohn, *Spin geometry*, Princeton Mathematical Series, 38, Princeton University Press, Princeton, NJ, 1989. MR1031992
- [32] N.C. Leung, “Riemannian geometry over different normed division algebras”, *J. Differential Geom.* **61** (2002), 289–333. MR1972148
- [33] J.-H. Lee and N.C. Leung, “Instantons and branes in manifolds with vector cross products”, *Asian J. Math.* **12** (2008), 121–143. MR2415016
- [34] K.F. Chan and N.C. Leung, “Calibrated Submanifolds in G_2 geometry”, *Lectures and Surveys on G_2 -manifolds and related topics*, Fields Institute Communications, Springer. (The present volume.)
- [35] J.D. Lotay, “Calibrated Submanifolds”, *Lectures and Surveys on G_2 -manifolds and related topics*, Fields Institute Communications, Springer. (The present volume.)
- [36] J.D. Lotay, “Geometric Flows of G_2 Structures”, *Lectures and Surveys on G_2 -manifolds and related topics*, Fields Institute Communications, Springer. (The present volume.)
- [37] W.S. Massey, “Obstructions to the existence of almost complex structures”, *Bull. Amer. Math. Soc.* **67** (1961), 559–564. MR0133137
- [38] J.W. Milnor and J D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, NJ, 1974. MR0440554
- [39] S. Salamon, *Riemannian geometry and holonomy groups*, Pitman Research Notes in Mathematics Series, 201, Longman Scientific & Technical, Harlow, 1989. MR1004008
- [40] D.A. Salamon and T. Walpuski, Notes on the octonions, in *Proceedings of the Gökova Geometry-Topology Conference 2016*, 1–85, Gökova Geometry/Topology Conference (GGT), Gökova. MR3676083
- [41] T. Walpuski, “Introduction to G_2 gauge theory”, *Lectures and Surveys on G_2 -manifolds and related topics*, Fields Institute Communications, Springer. (The present volume.)